

HOMOLOGICAL ALGEBRA FOR SCHWARTZ ALGEBRAS OF REDUCTIVE P-ADIC GROUPS

RALF MEYER

ABSTRACT. Let G be a reductive group over a non-Archimedean local field. Then the canonical functor from the derived category of smooth tempered representations of G to the derived category of all smooth representations of G is fully faithful. Here we consider representations on bornological vector spaces. As a consequence, if G is semi-simple, V and W are tempered irreducible representations of G , and V or W is square-integrable, then $\mathrm{Ext}_G^n(V, W) \cong 0$ for all $n \geq 1$. We use this to prove in full generality a formula for the formal dimension of square-integrable representations due to Schneider and Stuhler.

1. INTRODUCTION

Let G be a linear algebraic group over a non-Archimedean local field whose connected component of the identity element is reductive; we briefly call such groups *reductive p-adic groups*. For the purposes of exposition, we assume throughout the introduction that the connected centre of G is trivial, although we treat groups with arbitrary centre in the main body of this article.

We are going to compare homological and cohomological computations for the Hecke algebra $\mathcal{H}(G)$ and the Harish-Chandra Schwartz algebra $\mathcal{S}(G)$. Our main result asserts that the derived category of $\mathcal{S}(G)$ is a full subcategory of the derived category of $\mathcal{H}(G)$. These derived categories incorporate a certain amount of functional analysis because $\mathcal{S}(G)$ is more than just an algebra. Before we discuss this, we sketch two purely algebraic applications of our main theorem.

Let $\mathrm{Mod}_{\mathrm{alg}}(G)$ be the category of smooth representations of G on complex vector spaces. We compute some extension spaces in this Abelian category. If both V and W are irreducible tempered representations and one of them is square-integrable, then $\mathrm{Ext}_G^n(V, W) = 0$ for $n \geq 1$. If the local field underlying G has characteristic 0, this is proven by very different means in [20]. We get a more transparent proof that also works in prime characteristic.

The vanishing of $\mathrm{Ext}_G^n(V, W)$ is almost trivial if V or W is supercuspidal because then V or W is both projective and injective in $\mathrm{Mod}_{\mathrm{alg}}(G)$. This is related to the fact that supercuspidal representations are isolated points in the admissible dual. Square-integrable representations are isolated points in the tempered dual. Hence they are projective and injective in an appropriate category $\mathrm{Mod}(\mathcal{S}(G))$ of tempered smooth representations of G . Both $\mathrm{Mod}(\mathcal{S}(G))$ and $\mathrm{Mod}_{\mathrm{alg}}(G)$ are full subcategories in a larger category $\mathrm{Mod}(G)$. That is, we have fully faithful functors

$$\mathrm{Mod}(\mathcal{S}(G)) \rightarrow \mathrm{Mod}(G) \leftarrow \mathrm{Mod}_{\mathrm{alg}}(G).$$

We will show that the induced functors between the derived categories,

$$\mathrm{Der}(\mathcal{S}(G)) \rightarrow \mathrm{Der}(G) \leftarrow \mathrm{Der}_{\mathrm{alg}}(G),$$

are still fully faithful. This contains the vanishing result for Ext as a special case.

2000 *Mathematics Subject Classification*. 20G05, 18E30.

This research was supported by the EU-Network *Quantum Spaces and Noncommutative Geometry* (Contract HPRN-CT-2002-00280) and the *Deutsche Forschungsgemeinschaft* (SFB 478).

Another application involves Euler characteristics for square-integrable representations. Let V be an irreducible square-integrable representation of G . By a theorem of Joseph Bernstein, any finitely generated smooth representation of G , such as V , has a finite type projective resolution $P_\bullet \rightarrow V$. Its Euler characteristic is defined as

$$\chi(V) := \sum (-1)^n [P_n] \in K_0(\mathcal{H}(G)).$$

Since V is square-integrable, it is a projective $\mathcal{S}(G)$ -module and therefore defines a class $[V] \in K_0(\mathcal{S}(G))$. We show that the map $K_0(\mathcal{H}(G)) \rightarrow K_0(\mathcal{S}(G))$ induced by the embedding $\mathcal{H}(G) \rightarrow \mathcal{S}(G)$ maps $\chi(V)$ to $[V]$. This is useful because in [20] Peter Schneider and Ulrich Stuhler construct very explicit finite type projective resolutions, so that we get a nice formula for $[V] \in K_0(\mathcal{S}(G))$. This implies an explicit formula for the formal dimension of V , which is proven in [20] if V is supercuspidal or if G has characteristic 0. One consequence of this formula is that the formal dimensions are quantised, that is, they are all multiples of some $\alpha > 0$. This allows to estimate the number of irreducible square-integrable representations that contain a given representation of a compact open subgroup of G .

Although these applications can be stated purely algebraically, their proofs require functional analysis. We may view $\mathcal{S}(G)$ just as an algebra and consider the category $\mathbf{Mod}_{\text{alg}}(\mathcal{S}(G))$ of modules over $\mathcal{S}(G)$ in the algebraic sense as in [21]. However, the functor $\mathbf{Der}_{\text{alg}}(\mathcal{S}(G)) \rightarrow \mathbf{Der}_{\text{alg}}(G)$ fails to be fully faithful. This problem already occurs for $G = \mathbb{Z}$. The issue is that the tensor product of $\mathcal{S}(G)$ with itself plays a crucial role. If we work in $\mathbf{Mod}_{\text{alg}}(\mathcal{S}(G))$, we have to deal with $\mathcal{S}(G) \otimes \mathcal{S}(G)$, which appears quite intractable. In $\mathbf{Mod}(\mathcal{S}(G))$ we meet instead the much simpler completion $\mathcal{S}(G \times G)$ of this space.

Now it is time to explain briefly how we do analysis. I am an advocate of bornologies as opposed to topologies. This means working with bounded subsets and bounded maps instead of open subsets and continuous maps. General bornological vector spaces behave better than general topological vector spaces for purposes of representation theory and homological algebra (see [15, 17]). The spaces that we shall use here carry both a bornology and a topology, and both structures determine each other. Therefore, readers who are familiar with topological vector spaces may be able to follow this article without learning much about bornologies. We explain some notions of bornological analysis along the way because they may be unfamiliar to many readers.

We let $\mathbf{Mod}(G)$ be the category of smooth representations of G on bornological vector spaces as in [15]. The algebras $\mathcal{H}(G)$ and $\mathcal{S}(G)$ are bornological algebras in a natural way. A smooth representation $\pi: G \rightarrow \text{Aut}(V)$ on a bornological vector space V is called tempered if its integrated form extends to a bounded algebra homomorphism $\mathcal{S}(G) \rightarrow \text{End}(V)$. We may identify $\mathbf{Mod}(G)$ with the category of essential (or non-degenerate) bornological left modules over $\mathcal{H}(G)$ ([15]). As our notation suggests, this identifies the subcategory $\mathbf{Mod}(\mathcal{S}(G))$ with the category of essential bornological modules over $\mathcal{S}(G)$. We turn $\mathbf{Mod}(G)$ and $\mathbf{Mod}(\mathcal{S}(G))$ into exact categories using the class of extensions with a bounded linear section. The exact category structure allows us to form the derived categories $\mathbf{Der}(G)$ and $\mathbf{Der}(\mathcal{S}(G))$ as in [11]. Actually, the passage to derived categories is rather easy in both cases because our categories have enough projective and injective objects.

Equipping a vector space with the finest possible bornology, we identify the category of vector spaces with a full subcategory of the category of bornological vector spaces. Thus $\mathbf{Mod}_{\text{alg}}(G)$ becomes a full subcategory of $\mathbf{Mod}(G)$. Moreover, this embedding maps projective objects again to projective objects. Therefore, the induced functor $\mathbf{Der}_{\text{alg}}(G) \rightarrow \mathbf{Der}(G)$ is still fully faithful. Our main theorem asserts

that the canonical functor $\mathrm{Der}(\mathcal{S}(G)) \rightarrow \mathrm{Der}(G)$ is fully faithful as well. The basic technology for its proof is already contained in [16, 17].

In [17], I define the category $\mathrm{Mod}(A)$ of essential modules and its derived category $\mathrm{Der}(A)$ for a “quasi-unital” bornological algebra A and extend some homological machinery to this setting. A morphism $A \rightarrow B$ is called *isocohomological* if the induced functor $\mathrm{Der}(B) \rightarrow \mathrm{Der}(A)$ is fully faithful. [17] gives several equivalent characterisations of isocohomological morphisms. The criterion that is most easy to verify is the following: let $P_\bullet \rightarrow A$ be a projective A -bimodule resolution of A ; then $A \rightarrow B$ is isocohomological if and only if $B \hat{\otimes}_A P_\bullet \hat{\otimes}_A B$ is a resolution of B (in both cases, resolution means that there is a bounded contracting homotopy).

The article [16] deals with the special case of the embedding $\mathbb{C}[G] \rightarrow \mathcal{S}_1(G)$ for a finitely generated discrete group G and a certain Schwartz algebra $\mathcal{S}_1(G)$, which is defined by ℓ_1 -estimates. The chain complex whose contractibility decides whether this embedding is isocohomological turns out to be a *coarse geometric* invariant of G . That is, it depends only on the quasi-isometry class of a word-length function on G . If the group G admits a sufficiently nice combing, then I construct an explicit contracting homotopy of this chain complex. Thus $\mathbb{C}[G] \rightarrow \mathcal{S}_1(G)$ is isocohomological for such groups.

The argument for Schwartz algebras of reductive p -adic groups follows the same pattern. Let $H \subseteq G$ be some compact open subgroup and let $X := G/H$. This is a discrete space which inherits a canonical coarse geometric structure from G . Since G is reductive, it acts properly and cocompactly on a Euclidean building, namely, its affine Bruhat-Tits building. Such buildings are CAT(0) spaces and hence combable. Since X is coarsely equivalent to the building, it is combable as well. Thus the geometric condition of [16] is easily fulfilled for all reductive p -adic groups. However, we also have to check that the constructions in [16] are compatible with uniform smoothness of functions because G is no longer discrete. This forces us to look more carefully at the geometry of the building.

2. BORNOLOGICAL ANALYSIS

Algebras like the Schwartz algebra $\mathcal{S}(G)$ of a reductive p -adic group carry an additional structure that allows to do analysis in them. The homological algebra for modules over such algebras *simplifies* if we take this additional structure into account. One reason is that the complete tensor product $\mathcal{S}(G) \hat{\otimes} \mathcal{S}(G)$ can be identified with $\mathcal{S}(G^2)$ (Lemma 2).

It is customary to describe this additional structure using a locally convex topology. We prefer to use bornologies instead. This means that we work with bounded subsets and bounded operators instead of open subsets and continuous operators. A basic reference on bornologies is [9]. We use bornologies because of their advantages in connection with homological algebra (see [17]).

We mainly need bornological vector spaces that are complete and convex. Therefore, we drop these adjectives and tacitly require all bornologies to be complete and convex. When we use incomplete bornologies, we explicitly say so.

We need two classes of examples: fine bornologies and von Neumann bornologies. Let V be a vector space over $\mathbb{C}$. The *fine bornology* $\mathrm{Fine}(V)$ is the finest possible bornology on V . A subset $T \subseteq V$ is bounded in $\mathrm{Fine}(V)$ if and only if there is a finite-dimensional subspace $V_T \subseteq V$ such that T is a bounded subset of $V_T \cong \mathbb{R}^n$ in the usual sense. We also write $\mathrm{Fine}(V)$ for V equipped with the fine bornology.

Any linear map $\mathrm{Fine}(V) \rightarrow W$ is bounded. This means that Fine is a fully faithful functor from the category of vector spaces to the category of bornological vector spaces that is left-adjoint to the forgetful functor in the opposite direction.

Let V be a (quasi)complete locally convex topological vector space. A subset $T \subseteq V$ is called *von Neumann bounded* if it is absorbed by all neighbourhoods of zero. These subsets form a bornology on V called the *von Neumann bornology* (following [9]). We write $\mathbf{vN}(V)$ for V equipped with this bornology.

This defines a functor $\mathbf{vN}$ from topological to bornological vector spaces. Its restriction to the full subcategory of Fréchet spaces or, more generally, of LF-spaces, is fully faithful. That is, a linear map between such spaces is bounded if and only if it is continuous. A crucial advantage of bornologies is that joint boundedness is much weaker than joint continuity for multilinear maps: if $V_1, \dots, V_n, W$ are (quasi)complete locally convex topological vector spaces, then any separately continuous n -linear map $V_1 \times \dots \times V_n \rightarrow W$ is (jointly) bounded. The converse also holds under mild hypotheses.

Let G be a reductive p -adic group. We carefully explain how the Schwartz algebra $\mathcal{S}(G)$ looks like as a bornological algebra. The most convenient definition for our purposes is due to Marie-France Vignéras ([25]). Let $\sigma: G \rightarrow \mathbb{N}$ be the usual scale on G . It can be defined using a representation of G . Let $L_2(G)$ be the Hilbert space of square-integrable functions with respect to some Haar measure on G . Let

$$L_2^\sigma(G) := \{f: G \rightarrow \mathbb{C} \mid f \cdot \sigma^k \in L_2(G) \text{ for all } k \in \mathbb{N}\}.$$

A subset $T \subseteq L_2^\sigma(G)$ is bounded if for all $k \in \mathbb{N}$ there exists a constant $C_k \in \mathbb{R}_+$ such that $\|f \cdot \sigma^k\|_{L_2(G)} \leq C_k$ for all $f \in T$. This is the von Neumann bornology with respect to the Fréchet topology on $L_2^\sigma(G)$ defined by the sequence of semi-norms

$$\|f\|_2^k := \|f \cdot \sigma^k\|_{L_2(G)}.$$

Let $\mathbf{CO}(G)$ be the set of compact open subgroups of G , ordered by inclusion. For $U \in \mathbf{CO}(G)$, let $\mathcal{S}(G//U) = L_2^\sigma(G//U)$ be the subspace of U -bi-invariant functions in $L_2^\sigma(G)$. We give $\mathcal{S}(G//U)$ the subspace bornology, that is, a subset is bounded if and only if it is bounded in $L_2^\sigma(G)$. Finally, we let

$$\mathcal{S}(G) := \varinjlim \mathcal{S}(G//U),$$

where U runs through $\mathbf{CO}(G)$. We equip $\mathcal{S}(G)$ with the direct-limit bornology. That is, a subset of $\mathcal{S}(G)$ is bounded if and only if it is a bounded subset of $\mathcal{S}(G//U)$ for some $U \in \mathbf{CO}(G)$. We may also characterise this bornology as the von Neumann bornology with respect to the direct-limit topology on $\mathcal{S}(G)$, using the well-known description of bounded subsets in LF-spaces (see [24, Proposition 14.6]).

Lemma 1 ([25]). *The definition of the Schwartz algebra above agrees with the one of Harish-Chandra in [22, 26].*

Proof. The first crucial point is that the space of *double* cosets $G//U$ —as opposed to the group G itself—has polynomial growth with respect to the scale σ . It suffices to check this for a good maximal compact subgroup K because the map $G//U \rightarrow G//K$ is finite-to-one. By the Iwasawa decomposition, the double cosets in $G//K$ can be parametrised by points in a maximal split torus. The scale on G restricts to a standard word-length function on this torus, so that we get the desired polynomial growth. As a result, there exists $d > 0$ such that $\sum_{x \in G//U} \sigma^{-d}(x)$ is bounded.

Moreover, we need the following relationship between the growth of the double cosets UxU and the Harish-Chandra spherical function Ξ : there are constants $C, r > 0$ such that

$$\text{vol}(UxU) \leq C\sigma(x)^r \cdot \Xi(x)^{-2}, \quad \Xi(x)^{-2} \leq C\sigma(x)^r \cdot \text{vol}(UxU).$$

This follows from Equation I.1.(5) and Lemma II.1.1 in [26]. Hence

$$\begin{aligned} \int_G |f(x)|^2 \sigma(x)^s dx &= \sum_{x \in G//U} |f(UxU)|^2 \sigma(x)^s \text{vol}(UxU) \\ &\leq \sum_{x \in G//U} |f(UxU)|^2 \Xi(x)^{-2} C \sigma(x)^{r+s} \\ &\leq \max_{x \in G} |f(x)|^2 \Xi(x)^{-2} \sigma(x)^{r+s+d} \sum_{y \in G//U} C \sigma^{-d}(y). \end{aligned}$$

A similar computation shows

$$\int_G |f(x)|^2 \sigma(x)^s dx \geq \max_{x \in G} |f(x)|^2 \Xi(x)^{-2} C^{-1} \sigma(x)^{s-r}.$$

Therefore, the sequences of semi-norms $\|f\sigma^s\|_2$ and $\|f\Xi^{-1}\sigma^s\|_\infty$ for $s \in \mathbb{N}$ are equivalent and define the same function space $\mathcal{S}(G//U)$. $\square$

Convolution defines a continuous bilinear map $\mathcal{S}(G//U) \times \mathcal{S}(G//U) \rightarrow \mathcal{S}(G//U)$ for any $U \in \text{CO}(G)$ by [26, Lemme III.6.1]. Since $\mathcal{S}(G//U)$ is a Fréchet space, boundedness and continuity of the convolution are equivalent. Since any bounded subset of $\mathcal{S}(G)$ is already contained in $\mathcal{S}(G//U)$ for some U , the convolution is a bounded bilinear map on $\mathcal{S}(G)$, so that $\mathcal{S}(G)$ is a bornological algebra. In contrast, the convolution on $\mathcal{S}(G)$ is only separately continuous.

Now we return to the general theory and define the Hom functor and the tensor product. Let $\text{Hom}(V, W)$ be the vector space of bounded linear maps $V \rightarrow W$. A subset T of $\text{Hom}(V, W)$ is bounded if and only if it is *equibounded*, that is, $\{f(v) \mid f \in T, v \in S\}$ is bounded for any bounded subset $S \subseteq V$. This bornology is automatically complete if W is.

The *complete projective bornological tensor product* $\hat{\otimes}$ is defined in [8] by the expected universal property: it is a bornological vector space $V \hat{\otimes} W$ together with a bounded bilinear map $b: V \times W \rightarrow V \hat{\otimes} W$ such that $l \mapsto l \circ b$ is a bijection between bounded linear maps $V \hat{\otimes} W \rightarrow X$ and bounded bilinear maps $V \times W \rightarrow X$. This tensor product enjoys many useful properties. It is commutative, associative, and commutes with direct limits. It satisfies the adjoint associativity relation

$$(1) \quad \text{Hom}(V \hat{\otimes} W, X) \cong \text{Hom}(V, \text{Hom}(W, X)).$$

Therefore, a bornological module over a bornological algebra A can be defined in three equivalent ways, using a bounded linear map $A \rightarrow \text{End}(V)$, a bounded bilinear map $A \times V \rightarrow V$, or a bounded linear map $A \hat{\otimes} V \rightarrow V$.

Let $\otimes$ be the usual tensor product of vector spaces. The fine bornology functor is compatible with tensor products; that is, the obvious map $V \otimes W \rightarrow V \hat{\otimes} W$ is a bornological isomorphism

$$(2) \quad \text{Fine}(V \otimes W) \cong \text{Fine}(V) \hat{\otimes} \text{Fine}(W)$$

for any two vector spaces V and W . More generally, if W is any bornological vector space, then the underlying vector space of $\text{Fine}(V) \hat{\otimes} W$ is equal to the purely algebraic tensor product $V \otimes W$. A subset $T \subseteq V \otimes W$ is bounded if and only if there is a finite-dimensional subspace $V_T \subseteq V$ such that T is contained in and bounded in $V_T \otimes W \cong \mathbb{R}^n \otimes W \cong W^n$. Here W^n carries the direct-sum bornology. The reason for this is that $\hat{\otimes}$ commutes with direct limits.

If V_1 and V_2 are Fréchet-Montel spaces, then we have a natural isomorphism

$$(3) \quad \text{vN}(V_1 \hat{\otimes}_\pi V_2) \cong \text{vN}(V_1) \hat{\otimes} \text{vN}(V_2),$$

where $\hat{\otimes}_\pi$ denotes the *complete projective topological tensor product* (see [7, 24]). This isomorphism is proven in [13, Appendix A.1.4], based on results of Alexander

Grothendieck. The Montel condition means that all von Neumann bounded subsets are precompact (equivalently, relatively compact).

Lemma 2. *Let G be a reductive p -adic group. Then $\mathcal{S}(G) \hat{\otimes} \mathcal{S}(G) \cong \mathcal{S}(G \times G)$.*

Proof. It is shown in [25] that $\mathcal{S}(G//U)$ is a nuclear Fréchet space for all $U \in \text{CO}(G)$; in fact, this follows easily from the proof of Lemma 1. Equip G^2 with the scale $\sigma(a, b) := \sigma(a)\sigma(b)$ for all $a, b \in G$. By definition, $\mathcal{S}(G^2) \cong \varinjlim \mathcal{S}(G^2//U^2)$. Since $\hat{\otimes}$ commutes with direct limits,

$$\mathcal{S}(G) \hat{\otimes} \mathcal{S}(G) \cong \varinjlim \mathcal{S}(G//U) \hat{\otimes} \mathcal{S}(G//U)$$

as well. It remains to prove $\mathcal{S}(G//U)^{\hat{\otimes}^2} \cong \mathcal{S}(G^2//U^2)$. Since these Fréchet spaces are nuclear, they are Montel spaces. Hence (3) allows us to replace $\hat{\otimes}$ by $\hat{\otimes}_\pi$. Now we merely have to recall the definition of nuclearity (see [7, 24]).

Let V and W be Fréchet spaces. The natural map $V \otimes W \rightarrow \text{Hom}(V', W)$ defines another topology on $V \otimes W$, which may be weaker than the projective tensor product topology. A Fréchet space is nuclear if and only if this topology coincides with the projective tensor product topology. Equivalently, there is only one topology on $V \otimes W$ for which the canonical maps $V \times W \rightarrow V \otimes W$ and $V \otimes W \rightarrow \text{Hom}(V', W)$ are continuous. It is clear that the subspace topology from $\mathcal{S}(G^2//U^2)$ on $\mathcal{S}(G//U) \otimes \mathcal{S}(G//U)$ has these two properties. Hence it agrees with the projective tensor product topology. Now the assertion follows because $\mathcal{S}(G//U)^{\otimes^2}$ is dense in $\mathcal{S}(G^2//U^2)$. $\square$

3. BASIC HOMOLOGICAL ALGEBRA OVER THE HECKE ALGEBRA

Throughout this section, G denotes a totally disconnected, locally compact group, H denotes a fixed compact open subgroup of G , and $X := G/H$.

Let V be a bornological vector space and let $\pi: G \rightarrow \text{Aut}(V)$ be a representation of G by bounded linear operators. The representation π is called *smooth* if for any bounded subset $T \subseteq V$ there exists an open subgroup $U \subseteq G$ such that $\pi(g)v = v$ for all $g \in U$, $v \in T$ (see [15]). For example, the left and right regular representations of G on $\mathcal{S}(G)$ are smooth. If V carries the fine bornology, the definition above is equivalent to the usual notion of a smooth representation on a vector space.

Let $\text{Mod}(G)$ be the category of smooth representations of G on bornological vector spaces; its morphisms are the G -equivariant bounded linear maps. Let $\text{Mod}_{\text{alg}}(G)$ be the category of smooth representations of G on $\mathbb{C}$ -vector spaces. The fine bornology functor identifies $\text{Mod}_{\text{alg}}(G)$ with a full subcategory of $\text{Mod}(G)$.

Let $\mathcal{H}(G)$ be the Hecke algebra of G ; its elements are the locally constant, compactly supported functions on G . The convolution is defined by

$$f_1 * f_2(g) = \int_G f_1(x) f_2(x^{-1}g) dx$$

for some left-invariant Haar measure dx ; we normalise it so that $\text{vol}(H) = 1$. We equip $\mathcal{H}(G)$ with the fine bornology, so that $\mathcal{H}(G) \in \text{Mod}_{\text{alg}}(G) \subseteq \text{Mod}(G)$. More generally, given any bornological vector space V , we let $\mathcal{H}(G, V) := \mathcal{H}(G) \hat{\otimes} V$. The underlying vector space of $\mathcal{H}(G, V)$ is just $\mathcal{H}(G) \otimes V$ because $\mathcal{H}(G)$ carries the fine bornology. Hence $\mathcal{H}(G, V)$ is the space of locally constant, compactly supported functions $G \rightarrow V$. The *left regular representation* λ and the *right regular representation* ρ of G on $\mathcal{H}(G, V)$ are defined by

$$\lambda_g f(x) := f(g^{-1}x), \quad \rho_g f(x) := f(xg)$$

as usual. They are both smooth.

Any continuous representation $\pi: G \rightarrow \text{Aut}(V)$ on a bornological vector space V can be integrated to a bounded algebra homomorphism $\mathcal{H}(G) \rightarrow \text{End}(V)$, which we again denote by π . By adjoint associativity, this corresponds to a map

$$\pi_*: \mathcal{H}(G, V) = \mathcal{H}(G) \hat{\otimes} V \rightarrow V, \quad f \mapsto \int_G \pi(g, f(g)) dg.$$

The map π_* is G -equivariant if G acts on $\mathcal{H}(G, V)$ by λ . By [15, Proposition 4.7], the representation π is smooth if and only if π_* is a bornological quotient map, that is, any bounded subset of V is of the form $\pi_*(T)$ for some bounded subset $T \subseteq \mathcal{H}(G, V)$. Even more, if π is smooth, then π_* has a bounded linear section. Namely, we can use

$$(4) \quad \sigma_H: V \rightarrow \mathcal{H}(G, V), \quad \sigma_H v(g) = \pi(g^{-1}, v) 1_H(g),$$

where 1_H denotes the characteristic function of H . Thus the category $\text{Mod}(G)$ becomes isomorphic to the category $\text{Mod}(\mathcal{H}(G))$ of essential modules over $\mathcal{H}(G)$ (see [15, Theorem 4.8]). The term “essential” is a synonym for “non-degenerate” that is not as widely used for other purposes.

Let Ext be the class of all extensions in $\text{Mod}(G)$ that have a bounded linear section. This turns $\text{Mod}(G)$ into an exact category in the sense of Daniel Quillen. Hence the usual machinery of homological algebra applies to $\text{Mod}(G)$: we can form a derived category $\text{Der}(G)$ and derived functors (see [11, 17]). The exact category $\text{Mod}(G)$ has enough projective and injective objects, so that the usual recipes for computing derived functors apply. We shall use the following standard projective resolution in $\text{Mod}(G)$, which already occurs in [16].

The homogeneous space $X := G/H$ is discrete because H is open in G . Let

$$(5) \quad X_n := \{(x_0, \dots, x_n) \in X^{n+1} \mid x_0 \neq x_1, \dots, x_{n-1} \neq x_n\}.$$

We equip $\mathbb{C}[X_n]$ with the fine bornology. We let G act diagonally on X_n and equip $\mathbb{C}[X_n]$ with the induced representation

$$g \cdot f(x_0, \dots, x_n) := f(g^{-1}x_0, \dots, g^{-1}x_n).$$

The stabilisers of points

$$\text{Stab}(x_0, \dots, x_n) = \bigcap_{j=0}^n x_j H x_j^{-1}$$

are compact open subgroups of G for each $(x_0, \dots, x_n) \in X_n$. Let $X'_n \subseteq X_n$ be a subset that contains exactly one representative from each orbit. We get

$$(6) \quad X_n = \coprod_{\xi \in X'_n} G/\text{Stab}(\xi), \quad \mathbb{C}[X_n] = \bigoplus_{\xi \in X'_n} \mathbb{C}[G/\text{Stab}(\xi)].$$

If $U \in \text{CO}(G)$, then we have a natural isomorphism

$$\text{Hom}_G(\mathbb{C}[G/U], V) \xrightarrow{\cong} \text{Fix}(U, V), \quad f \mapsto f(1_U).$$

Since U is compact, this is an exact functor of V , so that $\mathbb{C}[G/U]$ is a projective object of $\text{Mod}(G)$. Therefore, $\mathbb{C}[X_n]$ is projective by (6).

In the following, we view X_n as a subset of $\mathbb{C}[X_n]$ in the usual way. We let $(x_0, \dots, x_n) = 0$ if $x_j = x_{j+1}$ for some $j \in \{0, \dots, n-1\}$. Thus $(x_0, \dots, x_n) \in \mathbb{C}[X_n]$ is defined for all $(x_0, \dots, x_n) \in X^{n+1}$.

We define the boundary map $\delta = \delta_n: \mathbb{C}[X_{n+1}] \rightarrow \mathbb{C}[X_n]$ for $n \in \mathbb{N}$ by

$$\delta((x_0, \dots, x_{n+1})) := \sum_{j=0}^{n+1} (-1)^j \cdot (x_0, \dots, \widehat{x_j}, \dots, x_{n+1}),$$

where $\widehat{x_j}$ means that x_j is omitted. In terms of functions, we can write

$$(7) \quad \delta\phi(x_0, \dots, x_n) = \sum_{j=0}^{n+1} (-1)^j \sum_{y \in X} \phi(x_0, \dots, x_{j-1}, y, x_j, \dots, x_n).$$

The operators δ_n are G -equivariant for all $n \in \mathbb{N}$. We define the augmentation map $\alpha: \mathbb{C}[X_0] \rightarrow \mathbb{C}$ by $\alpha(x) = 1$ for all $x \in X_0 = X$. It is G -equivariant with respect to the trivial representation of G on $\mathbb{C}$. It is easy to see that $\delta^2 = 0$ and $\alpha \circ \delta_0 = 0$. Hence we get a chain complex

$$C_\bullet(X) := (\mathbb{C}[X_n], \delta_n)_{n \in \mathbb{N}}$$

over $\mathbb{C}$. We also form the reduced complex $\tilde{C}_\bullet(X)$, which has $\mathbb{C}[X_n]$ in degree $n \geq 1$ and $\ker(\alpha: \mathbb{C}[X] \rightarrow \mathbb{C})$ in degree 0. The complex $\tilde{C}_\bullet(X)$ is exact. Thus $C_\bullet(X) \rightarrow \mathbb{C}$ is a projective resolution of $\mathbb{C}$ in $\mathbf{Mod}_{\text{alg}}(G)$.

Next we define *bivariant co-invariant spaces*. For $V, W \in \mathbf{Mod}(G)$, let $V \hat{\otimes}_G W$ be the quotient of $V \hat{\otimes} W$ by the closed linear span of $v \otimes w - gv \otimes gw$ for $v \in V$, $w \in W$, $g \in G$. Thus $V \hat{\otimes}_G W$ is again a complete bornological vector space. By definition, we have

$$V \hat{\otimes}_G W \cong W \hat{\otimes}_G V \cong (V \hat{\otimes} W) \hat{\otimes}_G \mathbb{C},$$

where we equip $V \hat{\otimes} W$ with the diagonal representation and $\mathbb{C}$ with the trivial representation of G . If X is another bornological vector space, then we may identify $\text{Hom}(V \hat{\otimes}_G W, X)$ with the space of bounded bilinear maps $f: V \times W \rightarrow X$ that satisfy $f(gv, gw) = f(v, w)$ for all $g \in G$, $v \in V$, $w \in W$. This universal property characterises $V \hat{\otimes}_G W$ uniquely. It follows from the defining property of $\hat{\otimes}$.

There is an alternative description of $V \hat{\otimes}_G W$ in terms of $\mathcal{H}(G)$ -modules. Turn V into a right and W into a left bornological $\mathcal{H}(G)$ -module by

$$v * f := \int_G f(g) g^{-1} v dg, \quad f * w := \int_G f(g) gw dg$$

for all $f \in \mathcal{H}(G)$, $v \in V$, $w \in W$. Let $V \hat{\otimes}_{\mathcal{H}(G)} W$ be the quotient of $V \hat{\otimes} W$ by the closed linear span of $v * f \otimes w - v \otimes f * w$ for $v \in V$, $f \in \mathcal{H}(G)$, $w \in W$.

Lemma 3. $V \hat{\otimes}_G W = V \hat{\otimes}_{\mathcal{H}(G)} W$, that is, the elements $gv \otimes gw - v \otimes w$ and $v * f \otimes w - v \otimes f * w$ generate the same closed linear subspace.

Proof. We have to show $\text{Hom}(V \hat{\otimes}_G W, X) = \text{Hom}(V \hat{\otimes}_{\mathcal{H}(G)} W, X)$ for all bornological vector spaces X . By definition, $\text{Hom}(V \hat{\otimes}_G W, X)$ is the space of bounded bilinear maps $l: V \times W \rightarrow X$ that satisfy $l(g^{-1}v, w) = l(v, gw)$ for all $v \in V$, $w \in W$, $g \in G$. This implies $l(v * f, w) = l(v, f * w)$ for all $v \in V$, $w \in W$, $f \in \mathcal{H}(G)$. Conversely, suppose $l(v * f, w) = l(v, f * w)$. Then

$$\begin{aligned} l(gv, g \cdot (f * w)) &= l(gv, (\delta_g * f) * w) \\ &= l(gv * (\delta_g * f), w) = l((g^{-1} \cdot gv) * f, w) = l(v, f * w) \end{aligned}$$

for all $v \in V$, $w \in W$, $g \in G$, $f \in \mathcal{H}(G)$. This implies $l(gv, gw) = l(v, w)$ for all $v \in V$, $w \in W$, $g \in G$ because any $w \in W$ is fixed by some $U \in \text{CO}(G)$ and therefore of the form $\mu_U * w$, where μ_U is the normalised Haar measure of U . $\square$

Since $\hat{\otimes}_G$ is functorial in both variables, we can apply it to chain complexes. Especially, we get a chain complex of bornological vector spaces $V \hat{\otimes}_G C_\bullet(X)$. Since $C_\bullet(X)$ is a projective resolution of the trivial representation, we denote the chain homotopy type of $V \hat{\otimes}_G C_\bullet(X)$ by $V \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$. The homology vector spaces of $V \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$ may be denoted $\text{Tor}_n^G(V, \mathbb{C})$ or $\text{Tor}_n^{\mathcal{H}(G)}(V, \mathbb{C})$. However, this passage to homology forgets an important part of the structure, namely, the bornology. Therefore, it is better to work with $V \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$ instead.

If V and W are just vector spaces, we can identify $V \hat{\otimes}_G W$ with a purely algebraic construction. Let $V \otimes_G W$ be the quotient of $V \otimes W$ by the linear span of $v \otimes w - gv \otimes gw$ for $v \in V$, $w \in W$, $g \in G$. Then

$$\text{Fine}(V \otimes_G W) \cong \text{Fine}(V) \hat{\otimes}_G \text{Fine}(W).$$

This follows from (2) and the fact that any linear subspace of a fine bornological vector space is closed. Therefore, $\text{Fine}(V) \hat{\otimes}_G C_\bullet(X) \cong \text{Fine}(V \otimes_G C_\bullet(X))$, and $\text{Tor}_*^G(V, \mathbb{C})$ is the homology of the chain complex $V \otimes_G C_\bullet(X)$. In this case, passage to homology is harmless because $V \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$ carries the fine bornology; this implies that it is quasi-isomorphic to its homology viewed as a complex with vanishing boundary map.

Our next goal is to describe $V \hat{\otimes}_G C_\bullet(X)$ (Proposition 6). This requires some geometric preparations.

Definition 4. Given a finite subset $F \subseteq X$, we define the relation $\sim_F$ on X by

$$(8) \quad x \sim_F y \iff (x, y) \in \bigcup_{g \in G} g \cdot (\{H\} \times F) \iff x^{-1}y \in HFH.$$

Here we view $x^{-1}y \in G//H$ and $HFH \subseteq G//H$.

A subset $S \subseteq X_n$ is *controlled by F* if $x_i \sim_F x_j$ for all $(x_0, \dots, x_n) \in S$ and all $i, j \in \{0, \dots, n\}$. We call $S \subseteq X_n$ *controlled* if it is controlled by some finite F . Roughly speaking, this means that all entries of S are uniformly close.

A subset $S \subseteq X_n$ is controlled if and only if S is G -finite, that is, there is a finite subset $F \subseteq X_n$ such that $S \subseteq G \cdot F$. This alternative characterisation will be used frequently. Definition 4 emphasises a crucial link between the controlled support condition and geometric group theory.

A *coarse (geometric) structure* on a locally compact space such as X is a family of relations on X satisfying some natural axioms due to John Roe (see also [6]). The subrelations of the relations $\sim_F$ above define a coarse geometric structure on X in this sense. Since it is generated by G -invariant relations, it renders the action of G on X isometric. This property already characterises the coarse structure uniquely: whenever a locally compact group acts properly and cocompactly on a locally compact space, there is a unique coarse structure for which this action is isometric (see [6, Example 6]). Moreover, with this coarse structure, the space X is coarsely equivalent to G .

By definition, the notion of a controlled subset of X_n depends only on the coarse geometric structure of X . Thus the space of functions on X_n of controlled support only depends on the large scale geometry of X .

Although our main examples, reductive groups, are unimodular, we want to treat groups with non-trivial modular function as well. Therefore, we have to decorate several formulas with modular functions. We define the modular homomorphism $\Delta_G: G \rightarrow \mathbb{R}_{>0}$ by $\Delta_G(g)d(g^{-1}) = dg$ and $d(gh) = \Delta_G(h)dg$ for all $h \in G$.

Definition 5. Let $\mathcal{C}(X_n, V)^\Delta$ be the space of all maps $\phi: X_n \rightarrow V$ that have controlled support and satisfy the covariance condition

$$(9) \quad \phi(g\xi) = \Delta_G(g)^{-1} \pi(g, \phi(\xi))$$

for all $\xi \in X_n$, $g \in G$. A subset $T \subseteq \mathcal{C}(X_n, V)^\Delta$ is bounded if $\{\phi(\xi) \mid \phi \in T\}$ is bounded in V for all $\xi \in X_n$ and the supports of all $\phi \in T$ are controlled by the same finite subset $F \subseteq X$.

Proposition 6. For any $V \in \text{Mod}(G)$, there is a natural bornological isomorphism

$$V \hat{\otimes}_G \mathbb{C}[X_n] \cong \mathcal{C}(X_n, V)^\Delta.$$

The induced boundary map on $V \hat{\otimes}_G \mathbb{C}[X_n]$ corresponds to the boundary map

$$\delta = \delta_n : \mathcal{C}(X_{n+1}, V)^\Delta \rightarrow \mathcal{C}(X_n, V)^\Delta$$

defined by (7).

We denote the resulting chain complex $(\mathcal{C}(X_n, V)^\Delta, \delta_n)_{n \in \mathbb{N}}$ by $\mathcal{C}(X_\bullet, V)^\Delta$.

Proof. The bifunctor $\hat{\otimes}_G$ commutes with direct limits and in particular with direct sums. Hence (6) yields

$$(10) \quad V \hat{\otimes}_G \mathbb{C}[X_n] \cong \bigoplus_{\xi \in X'_n} V \hat{\otimes}_G \mathbb{C}[G\xi] \cong \bigoplus_{\xi \in X'_n} V \hat{\otimes}_G \mathbb{C}[G/\text{Stab}(\xi)].$$

Fix $\xi \in X'_n$ and let $\text{Map}(G \cdot \xi, V)^\Delta$ be the space of all maps from $G \cdot \xi$ to V that satisfy the covariance condition (9). We equip $\text{Map}(G \cdot \xi, V)^\Delta$ with the product bornology as in Definition 5. We claim that the map

$$I : V \hat{\otimes}_G \mathbb{C}[G\xi] \rightarrow \text{Map}(G\xi, V), \quad v \otimes \phi \mapsto [\eta \mapsto \int_G \pi(h^{-1}, v) \cdot \phi(h\eta) dh],$$

yields a bornological isomorphism

$$V \hat{\otimes}_G \mathbb{C}[G\xi] \cong \text{Map}_G(G\xi, V)^\Delta.$$

We check that I descends to $V \hat{\otimes}_G \mathbb{C}[G\xi]$ and maps into $\text{Map}(G\xi, V)^\Delta$:

$$\begin{aligned} I(gv \otimes g\phi)(\eta) &= \int_G \pi(h^{-1}, gv) \cdot g\phi(h\eta) dh = \int_G \pi(h^{-1}g, v) \cdot \phi(g^{-1}h\eta) dh \\ &= \int_G \pi(h^{-1}, v) \cdot \phi(h\eta) dh = I(v \otimes \phi), \end{aligned}$$

$$\begin{aligned} I(v \otimes \phi)(g\eta) &= \int_G \pi(h^{-1}, v) \cdot \phi(hg\eta) dh \\ &= \int_G \pi(gh^{-1}, v) \phi(h\eta) d(hg^{-1}) = \Delta_G(g^{-1})\pi(g, I(v \otimes \phi)(\eta)). \end{aligned}$$

Thus we get a well-defined map $V \hat{\otimes}_G \mathbb{C}[G\xi] \rightarrow \text{Map}(G\xi, V)^\Delta$. Evaluation at ξ defines a bornological isomorphism $\text{Map}(G\xi, V)^\Delta \cong \text{Fix}(\text{Stab}(\xi), V)$. We claim that the latter is isomorphic to $V \hat{\otimes}_G \mathbb{C}[G/\text{Stab}(\xi)]$. Since $\text{Stab}(\xi)$ is compact and open, the Haar measure $\mu_{\text{Stab}(\xi)}$ of $\text{Stab}(\xi)$ is an element of $\mathcal{H}(G)$. Convolution on the right with $\mu_{\text{Stab}(\xi)}$ is an idempotent left module homomorphism on $\mathcal{H}(G)$, whose range is $\mathbb{C}[G/\text{Stab}(\xi)]$. Since $V \hat{\otimes}_G \mathcal{H}(G) \cong V$ for all V , additivity implies that $V \hat{\otimes}_G \mathbb{C}[G/\text{Stab}(\xi)]$ is equal to the range of $\mu_{\text{Stab}(\xi)}$ on V , that is, to $\text{Fix}(\text{Stab}(\xi), V)$. Thus we obtain an isomorphism $V \hat{\otimes}_G \mathbb{C}[G\xi] \cong \text{Map}_G(G\xi, V)^\Delta$, which can easily be identified with the map I .

Recall that a subset of X_n is controlled if and only if it meets only finitely many G -orbits. Therefore, we get the counterpart $\mathcal{C}(X_n, V)^\Delta \cong \bigoplus_{\xi \in X'_n} \text{Map}(G\xi, V)^\Delta$ to (10). We can piece our isomorphisms on orbits together to an isomorphism

$$I : V \hat{\otimes}_G \mathbb{C}[X_n] \rightarrow \mathcal{C}(X_n, V)^\Delta, \quad v \otimes \phi \mapsto [\xi \mapsto \int_G \pi(g^{-1}, v) \phi(g\xi) dg].$$

A straightforward computation yields $\delta \circ I(v \otimes \phi) = I(v \otimes \delta\phi)$ for all $v \in V$, $\phi \in \mathbb{C}[X_n]$ with δ as in (7). Therefore, I intertwines $\text{id} \hat{\otimes}_G \delta$ and δ . $\square$

Now let $C_\bullet(X, V) := C_\bullet(X) \hat{\otimes} V$, equipped with the diagonal representation of G . Since $C_\bullet(X)$ carries the fine bornology, the underlying vector space of $\mathbb{C}[X_n] \hat{\otimes} V$ may be identified with the space of functions $X_n \rightarrow V$ with finite support.

Lemma 7. *The chain complex $C_\bullet(X, V)$ is a projective resolution of V in $\text{Mod}(G)$.*

Proof. The complex $C_\bullet(X, V)$ is exact because $\hat{\otimes}$ is exact on extensions with a bounded linear section. We have $\text{Hom}_G(\mathbb{C}[G/U] \hat{\otimes} V, W) \cong \text{Hom}_U(V, W)$ for any $U \in \text{CO}(G)$ and any smooth representation W . Since this is an exact functor of W , $\mathbb{C}[G/U] \hat{\otimes} V$ is projective. Equation (6) shows that $\mathbb{C}[X_n] \hat{\otimes} V$ is a direct sum of such representations and therefore projective as well. $\square$

We view $\mathcal{H}(G)$ as a bimodule over itself in the usual way, by convolution on the left and right. Since right convolution commutes with the left regular representation, the complex $C_\bullet(X, \mathcal{H}(G))$ is a complex of $\mathcal{H}(G)$ -bimodules. The same reasoning as in the proof of Lemma 7 shows that it is a projective $\mathcal{H}(G)$ -bimodule resolution of $\mathcal{H}(G)$.

For $V, W \in \text{Mod}(G)$, we let $\text{Hom}_G(V, W)$ be the space of bounded G -equivariant linear maps $V \rightarrow W$, equipped with the equibounded bornology. It agrees with the space $\text{Hom}_{\mathcal{H}(G)}(V, W)$ of bounded linear $\mathcal{H}(G)$ -module homomorphisms. We also apply the bifunctor Hom_G to chain complexes. In particular, we can plug in the projective resolution $C_\bullet(X, V)$ of Lemma 7. The homotopy type of the resulting cochain complex of bornological vector spaces $\text{Hom}_G(C_\bullet(X, V), W)$ is denoted by $\mathbb{R}\text{Hom}_G(V, W)$. Its n th cohomology vector space is $\text{Ext}_G^n(V, W)$. As with $V \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$, it is preferable to retain the cochain complex itself.

If V and W carry the fine bornology, then $C_\bullet(X, V) = C_\bullet(X) \otimes V$ with the fine bornology. Therefore, $\mathbb{R}\text{Hom}_G(V, W)$ is equal to the space of all G -equivariant linear maps $C_\bullet(X) \otimes V \rightarrow W$. Hence the Ext spaces above agree with the purely algebraic Ext spaces. In more fancy language, the embedding $\text{Mod}_{\text{alg}}(G) \rightarrow \text{Mod}(G)$ induces a fully faithful functor between the derived categories $\text{Der}_{\text{alg}}(G) \rightarrow \text{Der}(G)$. This allows us to apply results proven using analysis in a purely algebraic context.

4. ISOCOHOMOLOGICAL SMOOTH CONVOLUTION ALGEBRAS

We introduce a class of convolution algebras on totally disconnected, locally compact groups G . These have the technical properties that allow us to formulate the problem. Then we examine the notion of an isocohomological embedding and formulate a necessary and sufficient condition for $\mathcal{H}(G) \rightarrow \mathcal{T}(G)$ to be isocohomological. This criterion involves the contractibility of a certain bornological chain complex, which is quite close to the one that arises in [16].

4.1. Unconditional smooth convolution algebras with rapid decay. Let G be a totally disconnected, locally compact group. Let $\sigma: G \rightarrow \mathbb{R}_{\geq 1}$ be a scale with the following properties: $\sigma(ab) \leq \sigma(a)\sigma(b)$ and $\sigma(a) = \sigma(a^{-1})$; σ is U -bi-invariant for some $U \in \text{CO}(G)$; the map σ is proper, that is, the subsets

$$(11) \quad B_R(G) := \{g \in G \mid \sigma(g) \leq R\}$$

are compact for all $R \geq 1$. The usual scale on a reductive p -adic groups has these properties. If the group G is finitely generated and discrete, then $\sigma = 1 + \ell$ or 2^ℓ for a word-length function ℓ are good, inequivalent choices.

Let $U \in \text{CO}(G)$. Given sets S, S' of functions $G//U \rightarrow \mathbb{C}$ we say that S' *dominates* S if for any $\phi \in S$ there exists $\phi' \in S'$ with $|\phi'(g)| \geq |\phi(g)|$ for all $g \in G$.

Definition 8. Let $\mathcal{T}(G)$ be a bornological vector space of functions $\phi: G \rightarrow \mathbb{C}$. We call $\mathcal{T}(G)$ an *unconditional smooth convolution algebra of rapid decay* if it satisfies the following conditions:

- 8.1. $\mathcal{T}(G)$ contains $\mathcal{H}(G)$;
- 8.2. $\mathcal{H}(G)$ is dense in $\mathcal{T}(G)$;
- 8.3. the convolution extends to a bounded bilinear map $\mathcal{T}(G) \times \mathcal{T}(G) \rightarrow \mathcal{T}(G)$;
- 8.4. $\mathcal{T}(G) = \varinjlim \mathcal{T}(G//U)$ as bornological vector spaces, where U runs through $\text{CO}(G)$ and $\mathcal{T}(G//U)$ is the space of U -bi-invariant functions in $\mathcal{T}(G)$;

- 8.5. if a set of functions $G//U \rightarrow \mathbb{C}$ is dominated by a bounded subset of $\mathcal{T}(G//U)$, then it is itself a bounded subset of $\mathcal{T}(G//U)$;
 8.6. M_σ is a bounded linear operator on $\mathcal{T}(G)$.

The first four conditions define a *smooth convolution algebra*, the fifth condition means that the convolution algebra is *unconditional*, the last one means that it has *rapid decay*.

An example of such a convolution algebra is the Schwartz algebra of a reductive p -adic group.

Let $\mathcal{T}(G)$ be an unconditional smooth convolution algebra of rapid decay. A representation $\pi: G \rightarrow \text{Aut}(V)$ is called $\mathcal{T}(G)$ -tempered if its integrated form extends to a bounded algebra homomorphism $\mathcal{T}(G) \rightarrow \text{End}(V)$ or, equivalently, to a bounded bilinear map $\mathcal{T}(G) \times V \rightarrow V$. The density of $\mathcal{H}(G)$ in $\mathcal{T}(G)$ implies that this extension is unique once it exists. Furthermore, G -equivariant maps are $\mathcal{T}(G)$ -module homomorphisms. Since the subalgebras $\mathcal{T}(G//U)$ are unital, the algebra $\mathcal{T}(G)$ is “quasi-unital” in the notation of [17], so that the category $\text{Mod}(\mathcal{T}(G))$ of essential bornological left $\mathcal{T}(G)$ -modules is defined. This category is naturally isomorphic to the category of $\mathcal{T}(G)$ -tempered smooth representations of G (see [17]). Thus $\text{Mod}(\mathcal{T}(G))$ is a full subcategory of $\text{Mod}(G)$.

The following lemmas prove some technical properties of $\mathcal{T}(G)$ that are obvious in most examples, anyway. Define $P_R: \mathcal{T}(G) \rightarrow \mathcal{H}(G)$ by $P_R\phi(x) = \phi(x)$ for $x \in B_R(G)$ and $P_R\phi(x) = 0$ otherwise, with $B_R(G)$ as in (11).

Lemma 9. $\lim_{R \rightarrow \infty} P_R(\phi) = \phi$ uniformly for ϕ in a bounded subset of $\mathcal{T}(G)$.

Proof. If $T \subseteq \mathcal{T}(G)$ is bounded, then $T \subseteq \mathcal{T}(G//U)$ for some $U \in \text{CO}(G)$. Shrinking U further, we achieve that the scale σ is U -bi-invariant. We may further assume that $\phi' \in T$ whenever $\phi': G//U \rightarrow \mathbb{C}$ is dominated by some $\phi \in T$ because $\mathcal{T}(G)$ is unconditional. Since M_σ is bounded, the subset $M_\sigma(T) \subseteq \mathcal{T}(G//U)$ is bounded as well. For any $\phi \in T$, we have $|\phi - P_R\phi| \leq R^{-1}|M_\sigma\phi|$, so that $\phi - P_R\phi \in R^{-1}M_\sigma(T)$. This implies uniform convergence $P_R(\phi) \rightarrow \phi$ for $\phi \in T$. $\square$

In the following, we briefly write

$$\mathcal{T}(G^2) := \mathcal{T}(G) \hat{\otimes} \mathcal{T}(G).$$

Lemma 2 justifies this notation for Schwartz algebras of reductive groups. In general, consider the bilinear maps

$$\mathcal{T}(G) \times \mathcal{T}(G) \rightarrow \mathbb{C}, \quad (\phi_1, \phi_2) \mapsto \phi_1(x)\phi_2(y)$$

for $(x, y) \in G^2$. They extend to bounded linear functionals on $\mathcal{T}(G^2)$ and hence map $\mathcal{T}(G^2)$ to a space of smooth functions on G^2 .

Lemma 10. *This representation of $\mathcal{T}(G^2)$ by functions on G^2 is faithful, that is, $\phi \in \mathcal{T}(G^2)$ vanishes once $\phi(x, y) = 0$ for all $x, y \in G$.*

Proof. The claim follows easily from Lemma 9 (this is a well-known argument in connection with Grothendieck’s Approximation Property). If $\phi(x, y) = 0$ for all $x, y \in G$, then also $(P_R \hat{\otimes} P_R)\phi(x, y) = 0$ for all $R \in \mathbb{N}$, $x, y \in G$. Since $P_R \hat{\otimes} P_R(\phi) \in \mathcal{H}(G^2)$, this implies $P_R \hat{\otimes} P_R(\phi) = 0$ for all $R \in \mathbb{N}$. Lemma 9 implies that $P_R \hat{\otimes} P_R$ converges towards the identity operator on $\mathcal{T}(G) \hat{\otimes} \mathcal{T}(G)$. This yields $\phi = 0$ as desired. $\square$

Hence we may view $\mathcal{T}(G^2)$ as a space of functions on G^2 . It is easy to see that $\mathcal{T}(G^2)$ is again a smooth convolution algebra on G^2 . Equip G^2 with the scale $\sigma_2(a, b) := \sigma(a)\sigma(b)$ for $a, b \in G$. Then the operator $M_{\sigma_2} = M_\sigma \hat{\otimes} M_\sigma$ is bounded, that is, $\mathcal{T}(G^2)$ also satisfies the rapid decay condition. However, $\mathcal{T}(G^2)$ need not

be unconditional. We *assume* $\mathcal{T}(G^2)$ to be *unconditional* in the following. This is needed for the proof of our main theorem.

Let $\mathcal{T}_c(G)$ be $\mathcal{H}(G)$ equipped with the subspace bornology from $\mathcal{T}(G)$. This bornology is incomplete, of course. Similarly, we let $\mathcal{T}_c(G^2)$ be $\mathcal{H}(G^2)$ equipped with the subspace bornology from $\mathcal{T}(G^2)$.

Lemma 11. *The completions of $\mathcal{T}_c(G)$ and $\mathcal{T}_c(G^2)$ are naturally isomorphic to $\mathcal{T}(G)$ and $\mathcal{T}(G^2)$.*

Proof. It suffices to prove this for $\mathcal{T}_c(G)$. We verify by hand that $\mathcal{T}(G)$ satisfies the universal property that defines the completion of $\mathcal{T}_c(G)$. Alternatively, we could use general characterisations of completions in [14, Section 4]. We must show that any bounded linear map $f: \mathcal{T}_c(G) \rightarrow W$ into a complete bornological vector space W extends uniquely to a bounded linear map on $\mathcal{T}(G)$. By Lemma 9, the sequence of operators $P_R: \mathcal{T}(G) \rightarrow \mathcal{T}_c(G)$ converges uniformly on bounded subsets towards the identity map on $\mathcal{T}(G)$. Hence any bounded extension $\bar{f}$ of f satisfies $\bar{f}(\phi) = \lim_{R \rightarrow \infty} f \circ P_R(\phi)$ for all $\phi \in \mathcal{T}(G)$. Conversely, this prescription defines a bounded linear extension of f . $\square$

4.2. Isocohomological convolution algebras. Let A be a quasi-unital algebra such as $\mathcal{H}(G)$ or $\mathcal{T}(G)$. In [17] I define the exact category $\mathbf{Mod}(A)$ of essential bornological left A -modules and its derived category $\mathbf{Der}(A)$. A bounded algebra homomorphism $f: A \rightarrow B$ between two quasi-unital bornological algebras induces functors $f^*: \mathbf{Mod}(B) \rightarrow \mathbf{Mod}(A)$ and $f^*: \mathbf{Der}(B) \rightarrow \mathbf{Der}(A)$. Trivially, if f has dense range then $f^*: \mathbf{Mod}(B) \rightarrow \mathbf{Mod}(A)$ is fully faithful. We call f *isocohomological* if $f^*: \mathbf{Der}(B) \rightarrow \mathbf{Der}(A)$ is fully faithful as well ([17]). We are interested in the embedding $\mathcal{H}(G) \rightarrow \mathcal{T}(G)$. If it is isocohomological, we briefly say that $\mathcal{T}(G)$ is *isocohomological*. The following conditions are proven in [17, Theorem 35] to be equivalent to $\mathcal{T}(G)$ being isocohomological:

- $V \hat{\otimes}_{\mathcal{T}(G)}^{\mathbb{L}} W \cong V \hat{\otimes}_G^{\mathbb{L}} W$ for all $V, W \in \mathbf{Der}(\mathcal{T}(G))$ (recall that $\hat{\otimes}_{\mathcal{H}(G)}^{\mathbb{L}} \cong \hat{\otimes}_G^{\mathbb{L}}$);
- $\mathbb{R} \operatorname{Hom}_{\mathcal{T}(G)}(V, W) \cong \mathbb{R} \operatorname{Hom}_G(V, W)$ for all $V, W \in \mathbf{Der}(\mathcal{T}(G))$;
- the functor $f^*: \mathbf{Der}(\mathcal{T}(G)) \rightarrow \mathbf{Der}(G)$ is fully faithful;
- $\mathcal{T}(G) \hat{\otimes}_G^{\mathbb{L}} V \cong V$ for all $V \in \mathbf{Mod}(\mathcal{T}(G))$;
- $\mathcal{T}(G) \hat{\otimes}_G^{\mathbb{L}} \mathcal{T}(G) \cong \mathcal{T}(G)$.

The last condition tends to be the easiest one to verify in practice. We will formulate it more concretely below. The signs “ $\cong$ ” in these statements mean isomorphism in the homotopy category of chain complexes of bornological vector spaces. This is stronger than an isomorphism of homology groups. As a consequence, we have $\operatorname{Tor}_n^G(V, W) \cong \operatorname{Tor}_n^{\mathcal{T}(G)}(V, W)$ and $\operatorname{Ext}_G^n(V, W) \cong \operatorname{Ext}_{\mathcal{T}(G)}^n(V, W)$ for all $V, W \in \mathbf{Mod}(\mathcal{T}(G))$ if $\mathcal{T}(G)$ is isocohomological.

Notions equivalent to that of an isocohomological embedding have been defined independently by several authors, as kindly pointed out to me by A. Yu. Pirkovskii (see [18] and the references given there). We warn the reader that in categories of topological algebras some of the conditions above are no longer equivalent. Namely, the cohomological conditions in terms of the derived category and $\mathbb{R} \operatorname{Hom}$ are weaker than the homological conditions involving $\hat{\otimes}^{\mathbb{L}}$.

We have seen in Section 3 that $V \hat{\otimes}_G W \cong (V \hat{\otimes} W) \hat{\otimes}_G \mathbb{C}$ for all $V, W \in \mathbf{Mod}(G)$, where $V \hat{\otimes} W$ is equipped with the diagonal representation of G . Since $V \hat{\otimes} W$ is projective if V or W is projective, this implies an isomorphism

$$V \hat{\otimes}_G^{\mathbb{L}} W \cong (V \hat{\otimes} W) \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$$

for all $V, W \in \text{Der}(G)$. Thus $\mathcal{T}(G)$ is isocohomological if and only if

$$(\mathcal{T}(G) \hat{\otimes} \mathcal{T}(G)) \hat{\otimes}_G^{\mathbb{L}} \mathbb{C} = \mathcal{T}(G^2) \hat{\otimes}_G^{\mathbb{L}} \mathbb{C} \cong \mathcal{T}(G).$$

Here we equip $\mathcal{T}(G^2)$ with the diagonal representation of G , which is given by

$$g \cdot f(x, y) := \Delta_G(g)f(xg, g^{-1}y)$$

for all $g \in G$, $f \in \mathcal{T}(G^2)$, $x, y \in G$ because the left and right $\mathcal{H}(G)$ -module structures on $\mathcal{T}(G)$ are the integrated forms of the left regular representation λ and the *twisted* right regular representation $\rho \cdot \Delta_G$. The convolution map

$$\mathcal{T}(G^2) = \mathcal{T}(G) \hat{\otimes} \mathcal{T}(G) \xrightarrow{*} \mathcal{T}(G)$$

descends to a bounded linear map $\mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C} \rightarrow \mathcal{T}(G)$. The latter map is a bornological isomorphism because $A \hat{\otimes}_A A \cong A$ for any quasi-unital bornological algebra by [17, Proposition 16]. Moreover, the convolution map $\mathcal{T}(G^2) \rightarrow \mathcal{T}(G)$ has a bounded linear section, namely, the map $\mathcal{T}(G) \rightarrow \mathcal{H}(G, \mathcal{T}(G)) \subseteq \mathcal{T}(G^2)$ defined in (4).

We may use the projective resolution $C_{\bullet}(X) \rightarrow \mathbb{C}$ to compute $\mathcal{T}(G^2) \hat{\otimes}_G^{\mathbb{L}} \mathbb{C}$. Proposition 6 identifies $\mathcal{T}(G^2) \hat{\otimes}_G C_{\bullet}(X)$ with $\mathcal{C}(X_{\bullet}, \mathcal{T}(G^2))^{\Delta}$. We augment this chain complex by the map

$$(12) \quad \alpha: \mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C}[X_0] \xrightarrow{\text{id} \hat{\otimes}_G \alpha} \mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C} \xrightarrow{*} \mathcal{T}(G)$$

We let $\tilde{\mathcal{C}}(X_{\bullet}, \mathcal{T}(G^2))^{\Delta}$ be the subcomplex of $\mathcal{C}(X_{\bullet}, \mathcal{T}(G^2))^{\Delta}$ that we get if we replace $\mathcal{C}(X_0, \mathcal{T}(G^2))^{\Delta}$ by $\tilde{\mathcal{C}}(X_0, \mathcal{T}(G^2))^{\Delta} := \ker \alpha$.

Proposition 12. *$\mathcal{T}(G)$ is isocohomological if and only if $\tilde{\mathcal{C}}(X_{\bullet}, \mathcal{T}(G^2))^{\Delta}$ has a bounded contracting homotopy.*

Proof. Our discussion of the convolution map implies that the augmentation map in (12) is a surjection with a bounded linear section. Hence it is a chain homotopy equivalence $\mathcal{T}(G^2) \hat{\otimes}_G^{\mathbb{L}} \mathbb{C} \rightarrow \mathcal{T}(G)$ if and only if its kernel $\tilde{\mathcal{C}}(X_0, \mathcal{T}(G^2))^{\Delta}$ is contractible. $\square$

To give the reader an idea why the various characterisations of isocohomological embeddings listed above are equivalent, we explain how the contractibility of $\tilde{\mathcal{C}}(X_{\bullet}, \mathcal{T}(G^2))^{\Delta}$ yields isomorphisms $\mathbb{R} \text{Hom}_{\mathcal{T}(G)}(V, W) \cong \mathbb{R} \text{Hom}_G(V, W)$ for $V, W \in \text{Mod}(\mathcal{T}(G))$. Almost the same argument yields $V \hat{\otimes}_{\mathcal{T}(G)}^{\mathbb{L}} W \cong V \hat{\otimes}_G^{\mathbb{L}} W$. The extension to objects of the derived categories is a mere formality.

The space $\mathcal{T}(G^2)$ carries a $\mathcal{T}(G)$ -bimodule structure via $f_1 * (f_2 \otimes f_3) * f_4 = (f_1 * f_2) \otimes (f_3 * f_4)$. This structure commutes with the inner conjugation action, so that $P_{\bullet} := \mathcal{T}(G^2) \hat{\otimes}_G C_{\bullet}(X)$ becomes a chain complex of bornological $\mathcal{T}(G)$ -bimodules over $\mathcal{T}(G)$. As above, we can compute these spaces explicitly:

$$\mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C}[X_n] \cong \bigoplus_{\xi \in X'_n} \text{Fix}(\text{Stab } \xi, \mathcal{T}(G^2)).$$

It is not hard to see that $\mathcal{T}(G^2)$ is a projective object of $\text{Mod}(\mathcal{T}(G^2))$. That is, $\mathcal{T}(G^2)$ is a projective bimodule. Since the summands of $\mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C}[X_n]$ are all retracts of $\mathcal{T}(G^2)$, we conclude that $\mathcal{T}(G^2) \hat{\otimes}_G \mathbb{C}[X_n]$ is a projective $\mathcal{T}(G)$ -bimodule.

Suppose now that $P_{\bullet}$ is a resolution of $\mathcal{T}(G)$. Then it is a projective $\mathcal{T}(G^2)$ -bimodule resolution. Since $\mathcal{T}(G)$ is projective as a right module, the contracting homotopy of $P_{\bullet}$ can be improved to consist of bounded right $\mathcal{T}(G)$ -module homomorphisms. Therefore, $P_{\bullet} \hat{\otimes}_{\mathcal{T}(G)} V$ is again a resolution of V . Explicitly,

$$P_n \hat{\otimes}_{\mathcal{T}(G)} V \cong \bigoplus_{\xi \in X'_n} \text{Fix}(\text{Stab}(\xi), \mathcal{T}(G) \hat{\otimes} V)$$

because $\mathcal{T}(G) \hat{\otimes}_{\mathcal{T}(G)} V \cong V$. The summands are retracts of the projective left $\mathcal{T}(G)$ -module $\mathcal{T}(G) \hat{\otimes} V$. Hence $P_\bullet \hat{\otimes}_{\mathcal{T}(G)} V$ is a projective left $\mathcal{T}(G)$ -module resolution of V . We use it to compute

$$\mathbb{R} \operatorname{Hom}_{\mathcal{T}(G)}(V, W) = \operatorname{Hom}_{\mathcal{T}(G)}(P_\bullet \hat{\otimes}_{\mathcal{T}(G)} V, W).$$

Let $U \in \operatorname{CO}(G)$ act on $\mathcal{T}(G) \hat{\otimes} V$ by $\Delta_G|_U \cdot \rho \otimes \pi = \rho \otimes \pi$. If $f: V \rightarrow W$ is bounded and U -equivariant, then $\phi \otimes v \mapsto \phi * f(v)$ defines a bounded G -equivariant linear map $\operatorname{Fix}(U, \mathcal{T}(G) \hat{\otimes} V) \rightarrow W$. One can show that this establishes a bornological isomorphism

$$\operatorname{Hom}_{\mathcal{T}(G)}(\operatorname{Fix}(U, \mathcal{T}(G) \hat{\otimes} V), W) \cong \operatorname{Hom}_U(V, W).$$

This yields a natural isomorphism

$$\operatorname{Hom}_{\mathcal{T}(G)}(P_n \hat{\otimes}_{\mathcal{T}(G)} V, W) \cong \bigoplus_{\xi \in X'_n} \operatorname{Hom}_{\operatorname{Stab}(\xi)}(V, W).$$

The right hand side no longer depends on $\mathcal{T}(G)$! Thus $\operatorname{Hom}_G(C_\bullet(X, V), W)$ is isomorphic to the same complex, and $\mathbb{R} \operatorname{Hom}_{\mathcal{T}(G)}(V, W) \cong \mathbb{R} \operatorname{Hom}_G(V, W)$ as asserted.

Next we simplify the chain complex $\mathcal{C}(X_\bullet, \mathcal{T}(G^2))^\Delta$. To $\phi \in \mathcal{C}(X_n, \mathcal{T}(G^2))^\Delta$ we associate a function $\phi_*: G \times G \times X_n \rightarrow \mathbb{C}$ by $\phi_*(g, h, \xi) := \phi(\xi)(g, h)$. This identifies $\mathcal{C}(X_n, \mathcal{T}(G^2))^\Delta$ with a space of functions on $G^2 \times X_n$ by Lemma 10. More precisely, we get the space of functions $\phi: G^2 \times X_n \rightarrow \mathbb{C}$ with the following properties:

- $\operatorname{supp} \phi \subseteq G^2 \times S$ for some controlled subset $S \subseteq X_n$;
- the function $(a, b) \mapsto \phi(a, b, \xi)$ belongs to $\mathcal{T}(G^2)$ for all $\xi \in X_n$;
- $\phi(ag, g^{-1}b, g^{-1}\xi) = \phi(a, b, \xi)$ for all $\xi \in X_n, g, a, b \in G$ (the two modular functions cancel).

The last condition means that ϕ is determined by its restriction to $\{1\} \times G \times X_n$ by $\phi(a, b, \xi) = \phi(1, ab, a\xi)$. Thus we identify $\mathcal{C}(X_n, \mathcal{T}(G^2))^\Delta$ with the following function space on $G \times X_n$:

Definition 13. Let $\mathcal{C}(G \times X_n, \mathcal{T})$ be the space of all functions $\phi: G \times X_n \rightarrow \mathbb{C}$ with the following properties:

- 13.1. $\operatorname{supp} \phi \subseteq G \times S$ for some controlled subset $S \subseteq X_n$;
- 13.2. the function $(a, b) \mapsto \phi(ab, a\xi)$ belongs to $\mathcal{T}(G^2)$ for all $\xi \in X_n$.

A subset $T \subseteq \mathcal{C}(G \times X_n, \mathcal{T})$ is bounded if there is a controlled subset $S \subseteq G$ such that $\operatorname{supp} \phi \subseteq G \times S$ for all $\phi \in T$ and if for any $\xi \in X_n$, the set of functions $(a, b) \mapsto \phi(ab, a\xi)$ for $\phi \in T$ is bounded in $\mathcal{T}(G^2)$.

The boundary map δ on $\mathcal{C}(X_n, \mathcal{T}(G^2))^\Delta$ corresponds to the boundary map

$$\begin{aligned} \delta: \mathcal{C}(G \times X_{n+1}, \mathcal{T}) &\rightarrow \mathcal{C}(G \times X_n, \mathcal{T}), \\ \delta\phi(g, x_0, \dots, x_n) &= \sum_{j=0}^{n+1} (-1)^j \sum_{y \in X} \phi(g, x_0, \dots, x_{j-1}, y, x_j, \dots, x_n). \end{aligned}$$

The augmentation map $\mathcal{C}(X, \mathcal{T}(G^2))^\Delta \rightarrow \mathcal{T}(G)$ corresponds to

$$(13) \quad \alpha: \mathcal{C}(G \times X, \mathcal{T}) \rightarrow \mathcal{T}(G), \quad \alpha\phi(g) = \sum_{x \in X} \phi(g, x).$$

The proofs are easy computations, which we omit. Let $\tilde{\mathcal{C}}(G \times X_0, \mathcal{T}) \subseteq \mathcal{C}(G \times X_0, \mathcal{T})$ be the kernel of α and let $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T})$ be the bornological chain complex that we get if we replace $\mathcal{C}(G \times X_0, \mathcal{T})$ by $\tilde{\mathcal{C}}(G \times X_0, \mathcal{T})$. Thus

$$\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}) \cong \tilde{\mathcal{C}}(X_\bullet, \mathcal{T}(G^2))^\Delta.$$

Smooth functions of compact support automatically satisfy both conditions in Definition 13, so that $\mathcal{H}(G) \otimes \mathbb{C}[X_n] \subseteq \mathcal{C}(G \times X_n, \mathcal{T})$. These embeddings are compatible with the boundary and augmentation maps. Thus $\mathcal{H}(G) \otimes \tilde{C}_\bullet(X)$ becomes a subcomplex of $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T})$. We write $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}_c)$ and $\mathcal{C}(G \times X_\bullet, \mathcal{T}_c)$ for the chain complexes $\mathcal{H}(G) \otimes \tilde{C}_\bullet(X)$ and $\mathcal{H}(G) \otimes C_\bullet(X)$ equipped with the incomplete subspace bornologies from $\mathcal{C}(G \times X_\bullet, \mathcal{T})$. The complex $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}_c)$ is contractible because $\tilde{C}_\bullet(X)$ is. However, the obvious contracting homotopy is unbounded.

Lemma 14. *Suppose that there is a contracting homotopy D for $\tilde{C}_\bullet(X)$ such that $\text{id}_{\mathcal{H}(G)} \otimes D$ is bounded on $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}_c)$. Then $\mathcal{T}(G)$ is isocohomological.*

Proof. We claim that $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T})$ is the completion of $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}_c)$. Then $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T}) \cong \tilde{\mathcal{C}}(X_\bullet, \mathcal{T}(G^2))^\Delta$ inherits a bounded contracting homotopy because completion is functorial. Proposition 12 yields that $\mathcal{T}(G)$ is isocohomological. It remains to prove the claim. We do this by reducing the assertion to Lemma 11. Since $\tilde{\mathcal{C}}(G \times X_\bullet, \mathcal{T})$ is a direct summand in $\mathcal{C}(G \times X_\bullet, \mathcal{T})$, it suffices to prove that $\mathcal{C}(G \times X_\bullet, \mathcal{T})$ is the completion of $\mathcal{C}(G \times X_\bullet, \mathcal{T}_c)$. Recall that $X'_\bullet$ denotes a subset of $X_\bullet$ containing one point from each G -orbit. The decomposition of $X_\bullet$ into G -orbits yields a direct-sum decomposition

$$(14) \quad \mathcal{C}(G \times X_\bullet, \mathcal{T}_c) \cong \bigoplus_{\xi \in X'_\bullet} \text{Fix}(\text{Stab } \xi, \mathcal{T}_c(G^2)),$$

and a similar decomposition for $\mathcal{C}(G \times X_\bullet, \mathcal{T})$. Here direct sums are equipped with the canonical bornology: a subset is bounded if it is contained in and bounded in a finite sub-sum. The reason for (14) is that a subset of X_n is controlled if and only if it meets only finitely many G -orbits. Since completion commutes with direct sums, the assertion now follows from Lemma 11. $\square$

5. CONTRACTING HOMOTOPIES CONSTRUCTED FROM COMBINGS

In order to apply Lemma 14, we have to construct contracting homotopies of $\tilde{C}_\bullet(X)$. For this we use the geometric recipes of [16]. The only ingredient is a sequence of maps $p_k: X \rightarrow X$ with certain properties. We first explain how such a sequence of maps gives rise to a contracting homotopy D of $\tilde{C}_\bullet(X)$. Then we formulate conditions on (p_k) and prove that they imply boundedness of D .

5.1. A recipe for contracting homotopies. The construction of $C_\bullet(X)$ and $\tilde{C}_\bullet(X)$ is natural: a map $f: X \rightarrow X$ induces a chain map $f_*: C_\bullet(X) \rightarrow C_\bullet(X)$ by $f_*((x_0, \dots, x_n)) := (f(x_0), \dots, f(x_n))$ or, equivalently,

$$(15) \quad f_*\phi(x_0, \dots, x_n) = \sum_{y_j: f(y_j)=x_j} \phi(y_0, \dots, y_n).$$

Since $\alpha \circ f_* = \alpha$, this restricts to a chain map on $\tilde{C}_\bullet(X)$. We have $\text{id}_* = \text{id}$ and $(fg)_* = f_*g_*$. Let p_0 be the constant map $x \mapsto H$ for all $x \in X$. We claim that $(p_0)_* = 0$ on $\tilde{C}_\bullet(X)$. On $\mathbb{C}[X_n]$ for $n \geq 1$ this is due to our convention that $(x_0, \dots, x_n) = 0$ if $x_i = x_{i+1}$ for some i . For $\phi \in \mathbb{C}[X_0]$, we get $(p_0)_*\phi = \alpha(\phi) \cdot (H)$, where $(H) \in \mathbb{C}[X]$ is the characteristic function of $H \in X$. This implies the claim.

Given maps $f, f': X \rightarrow X$, we define operators $D_j(f, f'): \mathbb{C}[X_n] \rightarrow \mathbb{C}[X_{n+1}]$ for $j \in \{0, \dots, n\}$ by

$$(16) \quad D_j(f, f')((x_0, \dots, x_n)) := (f(x_0), \dots, f(x_j), f'(x_j), \dots, f'(x_n))$$

and let $D(f, f') := \sum_{j=0}^n (-1)^j D_j(f, f')$. It is checked in [16] that

$$[\delta, D(f, f')] := \delta \circ D(f, f') + D(f, f') \circ \delta = f'_* - f_*.$$

Thus the chain maps f_* on $\tilde{C}_\bullet(X)$ for $f: X \rightarrow X$ are all chain homotopic. In particular, $D(\text{id}, p_0)$ is a contracting homotopy of $\tilde{C}_\bullet(X)$ because $(p_0)_* = 0$.

However, this trivial contracting homotopy does not work for Lemma 14. Instead, we use a sequence of maps $(p_k)_{k \in \mathbb{N}}$ with p_0 as above and $\lim_{k \rightarrow \infty} p_k = \text{id}$, that is, for each $x \in X$ there is $k_0 \in \mathbb{N}$ such that $p_k(x) = x$ for all $k \geq k_0$. We let

$$D_k := D(p_k, p_{k+1}), \quad D_{jk} := D_j(p_k, p_{k+1}).$$

Observe that D_{jk} vanishes on the basis vector $(x_0, \dots, x_n)$ unless $p_k(x_j) \neq p_{k+1}(x_j)$. Therefore, all but finitely many summands of

$$D := \sum_{k=0}^{\infty} D_k$$

vanish on any given basis vector. Thus D is a well-defined operator on $C_\bullet(X)$.

The operator D is a contracting homotopy of $\tilde{C}_\bullet(X)$ because

$$[D, \delta] = \sum_{k=0}^{\infty} [D(p_k, p_{k+1}), \delta] = \sum_{k=0}^{\infty} (p_{k+1})_* - (p_k)_* = \lim_{k \rightarrow \infty} (p_k)_* - (p_0)_* = \text{id}.$$

To verify this computation, plug in a basis vector and use that all but finitely many terms vanish. This is the operator we want to use in Lemma 14.

5.2. Sufficient conditions for boundedness. Construct D as above and let $D' := \text{id}_{\mathcal{H}(G)} \otimes D$. We want this to be a bounded operator on $\mathcal{C}(G \times X_\bullet, \mathcal{T}_c)$. For this, we impose three further conditions on (p_k) . First, (p_k) should be a *combing*. This notion comes from geometric group theory and is already used in [16]. It allows us to control the support of $D'\phi$ for ϕ with controlled support. Secondly, the combing (p_k) should be *smooth*. This allows us to control the smoothness of the functions $(a, b) \mapsto D'\phi(ab, a\xi)$ on G^2 for $\xi \in X_n$. Only the third condition involves the convolution algebra $\mathcal{T}(G)$. It asks for a certain sequence of operators $\mathcal{T}_c(G) \rightarrow \mathcal{T}_c(G^2)$ to be equibounded.

The smoothness condition is vacuous for discrete groups. The third condition is almost vacuous for ℓ_1 -Schwartz algebras of discrete groups. Hence these two conditions are not needed in [16]. In our application to reductive groups, we construct the operators (p_k) using the retraction of the affine Bruhat-Tits building of the group along geodesic paths. This is a combing because Euclidean buildings are CAT(0) spaces. Its smoothness amounts to the existence of congruence subgroups. The third condition follows easily from Lemma 2.

We now formulate the above conditions on (p_k) in detail and state the main result. We use the relation $\sim_F$ for a finite subset $F \subseteq X = G/H$ defined in (8) by $x \sim_F y \iff x^{-1}y \in HFH$.

Definition 15. A sequence of maps $(p_k)_{k \in \mathbb{N}}$ as above is called a *combing* of X if it has the following additional two properties:

- 15.1. there is a finite subset $F \subseteq X$ such that $p_k(x) \sim_F p_{k+1}(x)$ for all $k \in \mathbb{N}$, $x \in X$;
- 15.2. for any finite subset $F \subseteq X$ there is a finite subset $\bar{F} \subseteq X$ such that $p_k(x) \sim_{\bar{F}} p_k(y)$ for all $k \in \mathbb{N}$ and $x, y \in X$ with $x \sim_F y$.

We say that the combing has *polynomial growth* (with respect to the scale σ) if the least k_0 such that $p_k(gH) = gH$ for all $k \geq k_0$ grows at most polynomially in $\sigma(g)$. (This definition of growth differs slightly from the one in [16].)

We may view the sequence $(p_k(x))$ as a path from H to x . The conditions on a combing mean that these paths do not jump too far in each step and that nearby elements have nearby paths $(p_k(x))$.

Definition 16. A combing $(p_k)_{k \in \mathbb{N}}$ of G/H is called *smooth* if it has the following two properties:

- 16.1. all maps p_k are $\tilde{H}$ -equivariant for some open subgroup $\tilde{H} \subseteq H$;
- 16.2. for any $U \in \text{CO}(G)$, there exists $V \in \text{CO}(G)$ such that $aVb \subseteq UabU$ for all $a, b \in G$ with $p_k(abH) = aH$.

Definition 17. Let (p_k) be a smooth combing of polynomial growth. Define

$$R_k: \mathcal{T}_c(G) \rightarrow \mathcal{T}_c(G^2), \quad R_k \phi(a, b) := \begin{cases} \phi(ab) & \text{if } p_k(abH) = aH; \\ 0 & \text{otherwise.} \end{cases}$$

We say that (p_k) is *compatible with $\mathcal{T}(G)$* if the sequence of operators (R_k) is equibounded.

Theorem 18. *Let G be a totally disconnected, locally compact group and let $\mathcal{T}(G)$ be an unconditional smooth convolution algebra of rapid decay on G . Suppose also that the function space $\mathcal{T}(G^2)$ on G^2 is unconditional. If G/H for some compact open subgroup $H \subseteq G$ admits a smooth combing of polynomial growth that is compatible with $\mathcal{T}(G)$, then $\mathcal{T}(G)$ is isocohomological.*

5.3. Proof of Theorem 18.

Lemma 19. *Suppose that (p_k) is a combing. Then for any controlled subset $S \subseteq X_n$ there is a controlled subset $\tilde{S} \subseteq X_{n+1}$ such that $\text{supp } \phi \subseteq S$ implies $\text{supp } D(\phi) \subseteq \tilde{S}$ for all $\phi \in \mathbb{C}[X_n]$.*

Proof. Since S is controlled, there is a finite subset $F \subseteq X$ such that $x_i \sim_F x_j$ for all $i, j \in \{0, \dots, n\}$, $(x_i) \in S$. Since (p_k) is a combing, we can find a finite subset $\bar{F} \subseteq X$ such that $p_k(x_i) \sim_{\bar{F}} p_{k+1}(x_i)$ and $p_k(x_i) \sim_{\bar{F}} p_k(x_j)$ for all $k \in \mathbb{N}$ and all $(x_j) \in S$. Let $F' := H\bar{F}H\bar{F}H \subseteq G/H$. If $x \sim_{\bar{F}} y \sim_{\bar{F}} z$, then $x \sim_{F'} z$. Hence $(p_k(x_0), \dots, p_k(x_j), p_{k+1}(x_j), \dots, p_{k+1}(x_n))$ is controlled by F' for all $(x_i) \in S$. This means that all summands in $D(x_0, \dots, x_n)$ are controlled by F' . $\square$

Hence $D' := \text{id}_{\mathcal{H}(G)} \otimes D$ preserves controlled supports in $\mathcal{C}(G \times X_\bullet, \mathcal{T}_c)$. We have seen in (14) that

$$\mathcal{C}(G \times X_\bullet, \mathcal{T}_c) \cong \bigoplus_{\xi \in X'_\bullet} \text{Fix}(\text{Stab } \xi, \mathcal{T}_c(G^2)).$$

The isomorphism sends $\phi: G \times X_n \rightarrow \mathbb{C}$ to the family of functions $(\phi_\xi)_{\xi \in X'_n}$ defined by $\phi_\xi(a, b) := \phi(ab, a\xi)$. Thus we may describe any operator on $\mathcal{C}(G \times X_\bullet, \mathcal{T}_c)$ by a block matrix. In particular, we get $D' = (D'_{\xi\eta})_{\xi, \eta \in X'_\bullet}$ with certain operators

$$D'_{\xi\eta}: \text{Fix}(\text{Stab } \eta, \mathcal{T}_c(G^2)) \rightarrow \text{Fix}(\text{Stab } \xi, \mathcal{T}_c(G^2)).$$

The fact that D' preserves controlled supports means that for fixed η we have $D'_{\xi\eta} = 0$ for all but finitely many ξ . Thus the whole operator D' is bounded if and only if all its matrix entries $D'_{\xi\eta}$ are bounded.

For $j, n \in \mathbb{N}$, $n \geq j$, define $p_{jk}: X_n \rightarrow X^{n+2}$ by

$$p_{jk}((x_0, \dots, x_n)) := (p_k(x_0), \dots, p_k(x_j), p_{k+1}(x_j), \dots, p_{k+1}(x_n)).$$

Then the operator $D_{jk}: \mathbb{C}[X_n] \rightarrow \mathbb{C}[X_{n+1}]$ is given by

$$D_{jk}\phi(\xi) = \sum_{\eta \in p_{jk}^{-1}(\xi)} \phi(\eta).$$

Let $D'_{jk} = \text{id}_{\mathcal{H}(G)} \otimes D_{jk}$ and let $D'_{jk,\xi\eta}$ be the matrix entries of D'_{jk} with respect to the decomposition (14). Thus $D'_{\xi\eta} = \sum_{k \in \mathbb{N}} \sum_{j=0}^n (-1)^j D'_{jk,\xi\eta}$. Writing $\phi_\xi(a, b) = \phi(ab, a\xi)$ and $\phi_\eta(ag, g^{-1}b) = \phi(ab, ag\eta)$, we get

$$(17) \quad D'_{jk,\xi\eta}\psi(a, b) = \sum_{\{g \in G/\text{Stab}(\eta) \mid p_{jk}(ag\eta) = a\xi\}} \psi(ag, g^{-1}b) \\ = \text{vol}(\text{Stab} \eta)^{-1} \int_{\{g \in G \mid p_{jk}(ag\eta) = a\xi\}} \psi(ag, g^{-1}b) dg$$

for $\psi \in \text{Fix}(\text{Stab} \eta, \mathcal{T}_c(G^2))$. The right hand side of (17) makes sense for arbitrary $\psi \in \mathcal{T}_c(G^2)$ and extends $D'_{jk,\xi\eta}$ to an operator on $\mathcal{T}_c(G^2)$. Now we fix ξ, η until further notice and sometimes omit them from our notation.

Let $U \subseteq \text{Stab}(\eta)$ be an open subgroup and $\phi \in \mathcal{T}_c(G//U)$. Let $\mu_U \in \mathcal{H}(G)$ be the normalised Haar measure of U , that is, $\text{supp } \mu_U = U$ and $\mu_U(g) = \text{vol}(U)^{-1}$ for $g \in U$. Equation (17) yields

$$\text{vol}(\text{Stab} \eta) D'_{jk,\xi\eta}(\phi \otimes \mu_U)(a, b) \\ = \text{vol}(U)^{-1} \int_{\{g \in bU \mid p_{jk}(ag\eta) = a\xi\}} \phi(ag) dg = \begin{cases} \phi(ab) & \text{if } p_{jk}(ab\eta) = a\xi; \\ 0 & \text{otherwise.} \end{cases}$$

Let $\chi_{j,\xi\eta}(a, b)$ be the number of $k \in \mathbb{N}$ with $p_{jk}(ab\eta) = a\xi$ and let

$$\chi(a, b) = \chi_{\xi\eta}(a, b) := \sum_{j=0}^n (-1)^j \chi_{j,\xi\eta}(a, b).$$

These numbers are finite for any $a, b \in G$ for the same reason that guarantees that the sum defining D is finite on each basis vector. Define

$$A = A_{\xi\eta}: \mathcal{T}_c(G) \rightarrow \mathcal{T}_c(G^2), \quad A\phi(a, b) = \chi(a, b) \cdot \phi(ab).$$

Our computation shows that $A\phi = D'_{\xi\eta}(\phi \otimes \mu_U) \cdot \text{vol}(\text{Stab} \eta)$ if U is an open subgroup of $\text{Stab} \eta$ and $\phi \in \mathcal{T}_c(G//U)$.

Lemma 20. *The operator $D'_{\xi\eta}$ is bounded if and only if $A_{\xi\eta}$ is bounded.*

Proof. The boundedness of $D'_{\xi\eta}$ implies that A is bounded on $\mathcal{T}_c(G//U)$ for sufficiently small U and hence on all of $\mathcal{T}_c(G)$. Suppose conversely that A is bounded. We turn $\mathcal{T}_c(G^2)$ into an (incomplete) bornological right $\mathcal{T}_c(G)$ -module by

$$(f_1 \otimes f_2) * f_3 := f_1 \otimes (f_2 * f_3).$$

This bilinear map $\mathcal{T}_c(G^2) \times \mathcal{T}_c(G) \rightarrow \mathcal{T}_c(G^2)$ is bounded because the convolution in $\mathcal{T}(G)$ is bounded. The operators $D'_{jk,\xi\eta}$ and hence $D'_{\xi\eta}$ are $\mathcal{T}_c(G)$ -module homomorphisms by (17). Let $U \subseteq \text{Stab}(\eta)$ be open and $\phi_1, \phi_2 \in \mathcal{T}_c(G//U)$. Then

$$D'_{\xi\eta}(\phi_1 \otimes \phi_2) = D'_{\xi\eta}(\phi_1 \otimes \mu_U) * \phi_2 = \text{vol}(\text{Stab} \eta)^{-1} A(\phi_1) * \phi_2.$$

This implies the boundedness of $D'_{\xi\eta}$ because U is arbitrarily small and A and the convolution $\mathcal{T}_c(G^2) \rightarrow \mathcal{T}_c(G)$ are bounded. $\square$

Lemma 21. *If the combing is smooth, then for any $U \in \text{CO}(G)$ there is $V \in \text{CO}(G)$ such that A maps $\mathcal{T}_c(G//U)$ into $\mathcal{T}_c(G^2//V^2)$.*

Proof. Let $a, b \in G$. Clearly, $\chi(a, by) = \chi(a, b)$ for $y \in \text{Stab}(\eta)$. Since p_k is $\tilde{H}$ -equivariant, so is p_{jk} . Hence $\chi(ha, b) = \chi(a, b)$ for $h \in \tilde{H}$. Therefore, $A\phi(ua, b) = A\phi(a, b) = A\phi(a, bu)$ for $\phi \in \mathcal{T}_c(G//U)$ provided $U \subseteq \tilde{H} \cap \text{Stab}(\eta)$. Moreover, we have $A\phi(a, xb) = A\phi(ax, b)$ if $x \in \text{Stab}(\xi)$.

We may assume that the zeroth components η_0 and ξ_0 are H : any G -orbit on X_n has such a representative. Then $p_{jk}(ab\eta) = a\xi$ implies $p_k(abH) = aH$. By

the definition of a smooth combing, there is $V \in \mathbf{CO}(G)$ such that $aVb \subseteq UabU$ whenever $p_{jk}(ab\eta) = a\xi$ for some $k \in \mathbb{N}$. Hence $\phi(avb) = \phi(ab)$ for $(a, b) \in \text{supp } \chi$ and $v \in V$. We may shrink V such that $V \subseteq \text{Stab}(\xi) \cap U$. Then $\chi(a, vb) = \chi(a, b) = \chi(av, b)$ as well, so that A maps $\mathcal{T}_c(G//U)$ to $\mathcal{T}_c(G^2//V^2)$. $\square$

Definition 17 requires the following sequence of operators to be equibounded:

$$R_k : \mathcal{T}_c(G) \rightarrow \mathcal{T}_c(G^2), \quad R_k \phi(a, b) := \begin{cases} \phi(ab) & \text{if } p_k(abH) = aH; \\ 0 & \text{otherwise.} \end{cases}$$

Hence $R := \sum_{k \in \mathbb{N}} (k+1)^{-2} R_k$ is bounded. We have $R\phi(a, b) = \phi(ab)\chi'(a, b)$, where

$$\chi'(a, b) = \sum_{\{k \in \mathbb{N} \mid p_k(abH) = aH\}} (k+1)^{-2}.$$

Now let $S \subseteq \mathcal{T}_c(G)$ be bounded. Then $S \subseteq \mathcal{T}_c(G//U)$ for some $U \in \mathbf{CO}(G)$. We have already found $V \in \mathbf{CO}(G)$ such that $A\phi \in \mathcal{T}_c(G^2//V^2)$ for all $\phi \in \mathcal{T}_c(G//U)$. Since pointwise multiplication by the scale σ is bounded on $\mathcal{T}_c(G//V)$, it follows that the set of functions $\sigma(a)^N \sigma(b)^N R(S)$ is bounded in $\mathcal{T}_c(G^2//V^2)$. We claim that $\sigma(a)^N \sigma(b)^N R(S)$ dominates $A(S)$ for sufficiently large N . Since $\mathcal{T}(G^2)$ is unconditional, this implies the boundedness of A .

Since A and R are multiplication operators, the claim follows if $\chi' \sigma^N$ dominates $\chi_{j, \xi \eta}$ for any fixed j, ξ, η . This is what we are going to prove. Let $k_0(ab)$ be the least k_0 such that $p_k(ab\eta_j) = ab\eta_j$ for all $k \geq k_0$. The polynomial growth of the combing implies that $k_0(ab)$ is dominated by $C\sigma(ab)^N \leq C\sigma(a)^N \sigma(b)^N$ for sufficiently large C, N . Since $\xi_j \neq \xi_{j+1}$, we have $p_{jk}(ab\eta) \notin G\xi$ for $k \geq k_0$. Hence, if $p_{jk}(ab\eta) = a\xi$, then $k < k_0$. We choose the set of representatives $X'_\bullet$ such that $\xi_0 = H$ for all $\xi \in X'_\bullet$. Then $p_{jk}(ab\eta) = a\xi$ implies $p_k(abH) = aH$. Therefore, for each summand 1 in $\chi_{j, \xi \eta}(a, b)$ there is a summand $1/(k+1)^2 \geq 1/k_0^2$ in $\chi'(a, b)$. This yields the desired estimate $\chi_{j, \xi \eta}(a, b) \leq k_0(a, b)^2 \chi'(a, b) \leq C^2 \sigma(a)^{2N} \sigma(b)^{2N} \chi'(a, b)$. Thus the operators $A_{\xi \eta}$ are bounded for all ξ, η . This implies boundedness of $D'_{\xi \eta}$ by Lemma 20. By Lemma 19, it follows that D' is bounded. Finally, Lemma 14 yields that $\mathcal{T}(G)$ is isocohomological. This finishes the proof of Theorem 18.

6. A SMOOTH COMBING FOR REDUCTIVE p -ADIC GROUPS

The following theorem is the main goal of this section. In addition, we prove a variant (Theorem 28) that deals with the subcategories of χ -homogeneous representations for a character $\chi : C(G) \rightarrow \mathbf{U}(1)$ of the connected centre of G .

Theorem 22. *The Schwartz algebra $\mathcal{S}(G)$ of a reductive p -adic group G is isocohomological.*

Proof. We are going to apply Theorem 18. Let σ be the standard scale as in the definition of the Schwartz algebra and let $\mathcal{T}(G) := \mathcal{S}(G)$. The space $\mathcal{T}(G^2)$ is defined as $\mathcal{T}(G) \hat{\otimes} \mathcal{T}(G)$. This notation is permitted because of Lemma 2, which identifies $\mathcal{S}(G) \hat{\otimes} \mathcal{S}(G)$ with the Schwartz algebra of G^2 (which is again a reductive p -adic group). Clearly, the Schwartz algebras $\mathcal{S}(G)$ and $\mathcal{S}(G^2)$ are unconditional smooth convolution algebras of rapid decay.

We let $\mathcal{BT} = \mathcal{BT}(G)$ be the *affine Bruhat-Tits building* of G , as defined in [5, 23]. This is a Euclidean building on which G acts isometrically, properly, and cocompactly. Let $C(G)$ be the connected centre of G , so that the quotient $G/C(G)$ is semi-simple. In Section 7, we will also use the variant $\mathcal{BT}(G/C(G))$ of $\mathcal{BT}(G)$, which we call the *semi-simple affine Bruhat-Tits building* of G .

Let $G_\circ$ be the connected component of G as an algebraic group. Thus $G_\circ$ is a reductive group and G is a finite extension of $G_\circ$. Inspection of the definition in [23]

shows that the buildings for G and $G_\circ$ are equal. We remark that it is not hard to reduce the case of general reductive p -adic groups to the special case of connected semi-simple groups, or even connected simple groups. At first I followed this route myself. Eventually, it turned out that this intermediate step is unnecessary because all arguments work directly in the generality we need.

Let $\xi_0 \in \mathcal{BT}$, $H := \text{Stab}(\xi_0)$, and $X := G/H$. We have $H \in \text{CO}(G)$, and X may be identified with the discrete subset $G\xi_0 \subseteq \mathcal{BT}$. We need a combing of X . As a preparation, we construct a combing of $\mathcal{BT}$, using that Euclidean buildings are $\text{CAT}(0)$ spaces, that is, have “non-positive curvature” (see [4, 5]). In particular, any two points in $\mathcal{BT}$ are joined by a unique geodesic. For $\xi \in \mathcal{BT}$, let

$$p(\xi) : [0, d(\xi, \xi_0)] \rightarrow \mathcal{BT}, \quad t \mapsto p_t(\xi),$$

be the unit speed geodesic segment from ξ_0 to ξ ; extend this by $p_t(\xi) := \xi$ for $t > d(\xi, \xi_0)$. Restricting to $t \in \mathbb{N}$, we get a sequence of maps $p_k : \mathcal{BT} \rightarrow \mathcal{BT}$.

Lemma 23. *The maps p_k for $k \in \mathbb{N}$ form a combing of linear growth of $\mathcal{BT}$.*

This means $d(p_k(\xi), p_k(\eta)) \leq R \cdot d(\xi, \eta) + R$ and $d(p_k(\xi), p_{k+1}(\xi)) \leq R$ for all $\xi, \eta \in \mathcal{BT}$, $k \in \mathbb{N}$, for some $R > 0$. Linear growth means that the least k_0 such that $p_k(\xi)$ is constant for $k \geq k_0$ grows at most linearly in $d(\xi, \xi_0)$.

Proof. By construction, $d(p_s(\xi), p_t(\xi)) \leq |t - s|$ for all $s, t \in \mathbb{R}_+$, $\xi \in \mathcal{BT}$, and $p_s(\xi)$ is constant for $s \geq d(\xi, \xi_0)$. The lemma follows if we prove the following claim: $d(p_t(\xi), p_t(\eta)) \leq d(\xi, \eta)$ for all $\xi, \eta \in \mathcal{BT}$, $t \in \mathbb{R}_+$.

Fix $\xi, \eta \in \mathcal{BT}$ and $t \in \mathbb{R}_+$. We may assume $d(\xi, \xi_0) \geq d(\eta, \xi_0)$ (otherwise exchange ξ and η) and $d(\xi, \xi_0) \geq t$ (otherwise $p_t(\xi) = \xi$ and $p_t(\eta) = \eta$). Let d^* be the usual flat Euclidean metric on $\mathbb{R}^2$. Let ξ^* and η^* be points in $\mathbb{R}^2$ with

$$d^*(\xi^*, 0) = d(\xi, \xi_0), \quad d^*(\eta^*, 0) = d(\eta, \xi_0), \quad d^*(\xi^*, \eta^*) = d(\xi, \eta).$$

The $\text{CAT}(0)$ condition means that distances between points on the boundary of the geodesic triangle (ξ, η, ξ_0) are dominated by the distances between the corresponding points in the comparison triangle $(\xi^*, \eta^*, 0)$. Here the point $p_t(\xi)$ corresponds to the point $p_t(\xi)^*$ on $[0, \xi^*]$ of distance t from the origin. The point $p_t(\eta)$ corresponds to the point $p_t(\eta)^*$ of distance $\min\{t, d(\eta, \xi_0)\}$ from the origin. An easy




FIGURE 1. A comparison triangle

computation or a glance at Figure 1 shows $d^*(p_t(\xi)^*, p_t(\eta)^*) \leq d^*(\xi^*, \eta^*)$. By the $\text{CAT}(0)$ condition, this implies $d(p_t(\xi), p_t(\eta)) \leq d(\xi, \eta)$. $\square$

We identify G/H with the orbit $G\xi_0 \subseteq \mathcal{BT}$. Since the group action is cocompact, there is some $R > 0$ such that for any $\xi \in \mathcal{BT}$ there exists $\xi' \in G\xi_0$ with $d(\xi, \xi') < R$. We let $p'_k(\xi)$ for $\xi \in \mathcal{BT}$ be a point in $G\xi_0$ with $d(p'_k(\xi), p_k(\xi)) < R$. We claim that any such choice defines a combing of G/H .

If we equip $G\xi_0$ with the metric d from $\mathcal{BT}$, the maps p'_k on $G\xi_0$ still form a combing in the metric sense because they are “close” to the combing (p_k) . The

subspace metric from $\mathcal{BT}$ and the relations $\sim_F$ in Definition 15 generate the same coarse geometric structure on X . That is, for any $R > 0$ there is a finite subset $F \subseteq X$ such that $d(x, y) \leq R$ implies $x \sim_F y$, and for any finite subset $F \subseteq X$ there is $R > 0$ such that $x \sim_F y$ implies $d(x, y) \leq R$. This is easy to verify by hand. Alternatively, it follows from the uniqueness of coarse structures mentioned after Definition 15. Hence (p'_k) is a combing of G/H in the sense we need.

We also need the combing to be smooth. To get this, we must choose the base point ξ_0 and the approximations $p'_k(\xi)$ more carefully. This requires some geometric facts about the apartments in the building. Let K be the non-Archimedean local field over which G is defined. Let $S \subseteq G$ be a maximal K -split torus of G . We do not distinguish in our notation between the algebraic groups S and G and their locally compact groups of K -rational points. Let $X^*(S)$ and $X_*(S)$ be the groups of algebraic characters and cocharacters of S , respectively. The $\mathbb{R}$ -vector space $A := X_*(S) \otimes \mathbb{R}$ is the *basic apartment* of (G, S) .

Let $\Phi \subseteq X^*(S)$ be the set of roots of G relative to S . Choose a simple system of roots $\Delta \subseteq \Phi$ and let $A_+ \subseteq A$ be the corresponding closed Weyl chamber:

$$A_+ := \{x \in A \mid \alpha(x) \geq 0 \text{ for all } \alpha \in \Delta\}.$$

Let W be the *Weyl group* of the root system Φ . It is the Coxeter group generated by orthogonal reflections in the hyperplanes $\alpha(x) = 0$ for $\alpha \in \Delta$. The positive cone A_+ is a fundamental domain for this action, that is, $W(A_+) = A$ (see [10]).

Let $Z \subseteq G$ be the centraliser of S . There is a canonical homomorphism $\nu: Z \rightarrow A$ (see [23, (1.2)]). Its kernel is compact and its range is a lattice $\Lambda \subseteq A$; that is, Λ is a discrete and cocompact subgroup of A . Moreover, let $\Lambda_+ := \Lambda \cap A_+$. Since Λ is free Abelian, we can lift it to a subgroup of Z and view $\Lambda \subseteq Z \subseteq G$. Let Φ_{af} be the set of *affine roots* as in [23, (1.6)]. These are affine functions $\alpha: A \rightarrow \mathbb{R}$ of the form $\alpha(x) = \alpha_0(x) + \gamma$ with $\alpha_0 \in \Phi$ and certain $\gamma \in \mathbb{R}$. Recall that $\Phi \subseteq \Phi_{\text{af}}$ and that Φ_{af} is invariant under translation by Λ . The subsets of A of the form $\{x \in A \mid \alpha(x) = 0\}$ and $\{x \in A \mid \alpha(x) \geq 0\}$ for $\alpha \in \Phi_{\text{af}}$ are called *walls* and *half-apartments*, respectively.

We define the *closure* $\text{cl}(\Omega) \subseteq A$ of a non-empty subset $\Omega \subseteq A$ as the intersection of all closed half-apartments containing Ω (see [5, (7.1.2)]). We claim that

$$(18) \quad \text{cl}(\{0, \xi\}) = A_+ \cap (\xi - A_+)$$

for all $\xi \in \Lambda_+$. Let $B := A_+ \cap (\xi - A_+)$. This is an intersection of half-apartments containing 0 and ξ because $\Phi \cup (\xi - \Phi) \subseteq \Phi_{\text{af}}$. Hence $\text{cl}(\{0, \xi\}) \subseteq B$. It remains to show that any half-apartment C containing 0 and ξ also contains B . Let C be defined by the equation $\alpha \geq 0$ for some affine root α with linear part $\alpha_0 \in \Phi$. We distinguish the cases $\alpha_0 > 0$ and $\alpha_0 < 0$. If $\alpha_0 > 0$, then $\alpha(x) = \alpha_0(x) + \alpha(0) \geq \alpha_0(x)$ is non-negative on A_+ ; if $\alpha_0 < 0$, then $\alpha(\xi - x) = -\alpha_0(x) + \alpha(\xi) \geq -\alpha_0(x)$ is non-negative on A_+ , so that α is non-negative on $\xi - A_+$. Thus $B \subseteq C$ in either case. This finishes the proof that $\text{cl}(\{0, \xi\}) = B$.

Lemma 24. *There is $R > 0$ such that for all $\xi \in \Lambda_+$ and all $\eta \in \text{cl}(\{0, \xi\})$, there is $\eta' \in \Lambda \cap \text{cl}(\{0, \xi\})$ with $d(\eta, \eta') \leq R$.*

Proof. Let $\Delta = \{\alpha_1, \dots, \alpha_s\}$ be the system of simple roots that determines A_+ . If G is semi-simple, these roots form a basis of A . In general, they are linearly independent, so that we can extend them to a basis by certain α_j for $s < j \leq r$. Define a vector space isomorphism $\gamma: A \rightarrow \mathbb{R}^r$ by $\gamma(\eta)_j = \alpha_j(\eta)$ for $j = 1, \dots, r$. This identifies A_+ with the set of $(x_j) \in \mathbb{R}^r$ with $x_j \geq 0$ for $1 \leq j \leq s$. Equation (18) identifies $B := \text{cl}(\{0, \xi\})$ with

$$\gamma(B) = \{(x_j) \in \mathbb{R}^r \mid 0 \leq x_j \leq \gamma(\xi)_j, j = 1, \dots, s\}.$$

We may assume $\alpha_j \in X_*(S) \otimes \mathbb{Q} \subseteq A$, so that $\gamma(\Lambda) \subseteq \mathbb{Q}^r$. Replacing γ by $n^{-1}\gamma$ for some $n \in \mathbb{N}^*$, we can achieve $\mathbb{Z}^r \subseteq \gamma(\Lambda)$. Hence if $(x_j) \in \gamma(B)$, then the truncated vector $\lfloor (x_j) \rfloor := (\lfloor x_j \rfloor)$ belongs to $\gamma(B \cap \Lambda)$. It satisfies $|x_j - \lfloor x_j \rfloor| < 1$ for all j . Since the norm $\|\gamma(\eta)\|_\infty$ is equivalent to the Euclidean norm on A , we have $d(\gamma^{-1}\lfloor \gamma(\eta) \rfloor, \eta) < R$ for all $\eta \in B$ for some $R > 0$. $\square$

The building $\mathcal{BT}$ can be defined as the quotient of $G \times A$ by a certain equivalence relation. We may view gA for $g \in G$ as a subspace of $\mathcal{BT}$; these are the *apartments* of $\mathcal{BT}$. We now choose ξ_0 to be the origin in $A \subseteq \mathcal{BT}$. Recall that $H \subseteq G$ denotes the stabiliser of ξ_0 . We have the *Cartan decomposition* $G = H\Lambda_+H$ by [23, (3.3.3)], so that $G\xi_0 = H\Lambda_+\xi_0$. Let $G_\circ \subseteq G$ be the connected component of the identity (as an algebraic variety) and let $H_\circ := H \cap G_\circ$. These are open normal subgroups of finite index in G and H , respectively, and $\mathcal{BT}(G)$ is isomorphic to $\mathcal{BT}(G_\circ)$ equipped with a canonical action of G .

Choose $\Lambda'_+ \subseteq \Lambda_+$ to contain one representative for each H -orbit in $G\xi_0$. Fix $\xi \in \Lambda'_+$ and $k \in \mathbb{N}$. We further decompose $H\xi$ as a disjoint union of finitely many $H_\circ$ -orbits $H_\circ h_j \xi = h_j H_\circ \xi$ for suitable $h_1, \dots, h_N \in H$. We let $p'_k(\xi)$ be some point in $\Lambda \cap \text{cl}(\{0, \xi\}) \subseteq A \subseteq \mathcal{BT}$ that has minimal distance from $p_k(\xi)$.

Proposition 25. *Let $\Omega \subseteq A \subseteq \mathcal{BT}$, $\Omega \neq \emptyset$. If $g \in G_\circ$ satisfies $gx = x$ for all $x \in \Omega$, then $gx = x$ for all $x \in \text{cl}(\Omega)$. (This may fail if we allow $g \in G$.)*

Proof. The proof requires some facts about stabilisers of points in $\mathcal{BT}$, which are conveniently summarised in [20, Section I.1]. The subgroups $P_\Omega \subseteq G_\circ$ defined there manifestly satisfy $P_\Omega = P_{\text{cl}(\Omega)}$. This implies our claim because

$$P_\Omega = \{g \in G_\circ \mid gx = x \ \forall x \in \Omega\}. \quad \square$$

Proposition 25 allows us to define $p'_k(h_j h_\circ \xi) := h_j h_\circ p'_k(\xi)$ for all $h_\circ \in H_\circ$. Letting ξ, j vary, we get a map $p'_k: G\xi_0 \rightarrow G\xi_0$. It is $H_\circ$ -equivariant because $H_\circ$ is normal in H . Since $p_k(\xi) \in \text{cl}(\{0, \xi\})$, Lemma 24 yields $R > 0$ such that $d(p_k(\xi), p'_k(\xi)) \leq R$ for all $k \in \mathbb{N}$, $\xi \in \Lambda'_+$. The same holds for $\xi \in G\xi_0$ because p_k is H -equivariant and G acts isometrically on $\mathcal{BT}$. Moreover, $p'_0(\xi) = \xi_0$ and $p'_k(\xi) = \xi$ for $k \geq d(\xi, \xi_0)$. Thus (p'_k) is a combing of G/H of linear growth.

Lemma 26. *The combing (p'_k) is smooth.*

Proof. There is a decreasing sequence $(U_n)_{n \in \mathbb{N}}$ in $\text{CO}(G)$ such that each U_n is normal in H and can be written as $U_n^+ \cdot U_n^- = U_n^- \cdot U_n^+$ with

$$(19) \quad \lambda U_n^+ \subseteq U_n^+ \lambda, \quad U_n^- \lambda \subseteq \lambda U_n^-$$

for all $\lambda \in \Lambda_+$ (see [20, Section I.2]). Let $a, b \in G$ satisfy $p'_k(abH) = aH$. Write $ab = h_1 \lambda h_2$ with $h_1, h_2 \in H$, $\lambda \in \Lambda'_+$, and h_1 chosen as carefully as above if G is disconnected. Then $a = h_1 p'_k(\lambda) h_3$ with the same h_1 and some $h_3 \in H$. Hence $b = h_3^{-1} p'_k(\lambda)^{-1} \lambda h_2$. Since $p'_k(\lambda) \in \text{cl}(\{0, \lambda\})$, equation (18) yields $p'_k(\lambda) \in \Lambda_+$ and $p'_k(\lambda)^{-1} \lambda \in \Lambda_+$. Using (19) and that U_n is normal in H , we get

$$\begin{aligned} aU_n b &= h_1 p'_k(\lambda) U_n^+ U_n^- p'_k(\lambda)^{-1} \lambda h_2 \subseteq h_1 U_n^+ p'_k(\lambda) p'_k(\lambda)^{-1} \lambda U_n^- h_2 \\ &\subseteq h_1 U_n \lambda U_n h_2 = U_n h_1 \lambda h_2 U_n = U_n a b U_n. \end{aligned}$$

Thus the combing (p'_k) is smooth in the sense of Definition 16. $\square$

Finally, the compatibility condition of Definition 17 is easy to check using the explicit description of $\mathcal{S}(G^2)$ in Lemma 2. In order to cover also the Schwartz algebras for discrete groups, which are defined by ℓ_1 -estimates, we define spaces

$$L_p^\sigma(G) := \{f: G \rightarrow \mathbb{C} \mid f \cdot \sigma^k \in L_p(G) \ \forall k \in \mathbb{N}\}$$

for all $1 \leq p < \infty$, and equip them with the evident bornology: a subset $T \subseteq L_p^\sigma(G)$ is bounded if and only if for any $k \in \mathbb{N}$ there is C_k such that $\|f \cdot \sigma^k\|_{L_p(G)} \leq C_k$ for all $f \in T$. Let $L_p^\sigma(G//U)$ be the subspace of U -bi-invariant functions in $L_p^\sigma(G)$.

For $p = 2$, this agrees with the previous definition, so that Lemma 2 yields

$$\mathcal{S}(G) = \varinjlim L_2^\sigma(G//U), \quad \mathcal{S}(G^2) = \varinjlim L_2^\sigma(G^2//U^2).$$

Lemma 27. *Let (p_k) be a combing of polynomial growth on G/H . Then the sequence of operators (R_k) used in Definition 17 is uniformly bounded as operators $L_p^\sigma(G) \rightarrow L_p^\sigma(G^2)$. Here we use the scale $\sigma(a, b) := \sigma(a)\sigma(b)$ on G^2 .*

Proof. The operator $W\phi(x, y) := \phi(x, x^{-1}y)$ is an isometry of $L_p(G^2)$. It is also sufficiently compatible with the scale on G^2 for W and its inverse to be bounded linear operators on $L_p^\sigma(G^2)$. We have $R_k\phi(a, b) = \phi(ab)1_{p_k(abH)}(a)$, where $1_{p_k(abH)}$ denotes the characteristic function of $p_k(abH) \subseteq G$. Hence

$$WR_k\phi(x, y) = \phi(y) \cdot 1_{p_k(yH)}(x).$$

Since the combing (p_k) has polynomial growth, $\sigma(p_k(yH))$ is controlled by a polynomial in $\sigma(y)$. The boundedness of $W \circ R_k$ is now immediate because all cosets xH have volume 1. This implies the boundedness of R_k . $\square$

Since the combing (p'_k) is smooth, for any $U \in \mathbf{CO}(G)$ there exists $V \in \mathbf{CO}(G)$ such that R_k maps $\mathcal{T}_c(G//U)$ into $\mathcal{T}_c(G^2//V^2)$. Together with Lemma 27 for $p = 2$, this yields that the combing (p'_k) is compatible with $\mathcal{S}(G)$ in the sense of Definition 17. We have now verified all the hypotheses of Theorem 18. Thus $\mathcal{S}(G)$ is isocohomological. $\square$

6.1. Decomposition with respect to the centre of G . As before, we let G be a reductive p -adic group. Let $C(G) \subseteq G$ be the connected centre of G and let $\chi: C(G) \rightarrow \mathbf{U}(1)$ be a unitary character on $C(G)$. Let $\mathbf{Mod}_\chi(G)$ be the full subcategory of $\mathbf{Mod}(G)$ whose objects are the representations $\pi: G \rightarrow \mathbf{Aut}(V)$ that satisfy $\pi(z) = \chi(z)\mathrm{id}_V$ for all $z \in C(G)$. Let

$$\mathbf{Mod}_\chi(\mathcal{S}(G)) := \mathbf{Mod}(\mathcal{S}(G)) \cap \mathbf{Mod}_\chi(G)$$

be the subcategory of tempered representations in $\mathbf{Mod}_\chi(G)$. The class of extensions with a bounded linear section turns $\mathbf{Mod}_\chi(G)$ and $\mathbf{Mod}_\chi(\mathcal{S}(G))$ into exact categories, so that we can form the derived categories $\mathbf{Der}_\chi(G)$ and $\mathbf{Der}_\chi(\mathcal{S}(G))$. Let $G_{\mathrm{ss}} := G/C(G)$, this is again a reductive p -adic group. If $\chi = 1$, then $\mathbf{Mod}_\chi(G) = \mathbf{Mod}(G_{\mathrm{ss}})$ and $\mathbf{Mod}_\chi(\mathcal{S}(G)) = \mathbf{Mod}(\mathcal{S}(G_{\mathrm{ss}}))$. In general, there are quasi-unital bornological algebras $\mathcal{H}_\chi(G)$ and $\mathcal{S}_\chi(G)$ such that

$$\mathbf{Mod}_\chi(G) \cong \mathbf{Mod}(\mathcal{H}_\chi(G)), \quad \mathbf{Mod}_\chi(\mathcal{S}(G)) \cong \mathbf{Mod}(\mathcal{S}_\chi(G)).$$

We briefly recall their well-known definitions. A $C(G)$ -invariant subset of G is called $C(G)$ -compact if its image in G_{ss} is compact. Let $\mathcal{H}_\chi(G)$ be the space of locally constant functions $f: G \rightarrow \mathbb{C}$ with $C(G)$ -compact support such that $f(z^{-1}g) = \chi(z)f(g)$ for all $g \in G$, $z \in C(G)$. If $f_1, f_2 \in \mathcal{H}_\chi(G)$, then the function $h \mapsto f_1(h)f_2(h^{-1}g)$ is $C(G)$ -invariant, so that

$$f_1 * f_2(g) := \int_{G_{\mathrm{ss}}} f_1(h)f_2(h^{-1}g) dh$$

makes sense; here dh denotes the Haar measure on G_{ss} . This turns $\mathcal{H}_\chi(G)$ into an algebra, which we equip with the fine bornology.

Let $L_2^\sigma(G)_\chi$ be the space of functions $f: G \rightarrow \mathbb{C}$ that satisfy $f(z^{-1}g) = \chi(z)f(g)$ for all $g \in G$, $z \in C(G)$, and such that $zg \mapsto |f(g)|$ is an element of $L_2^\sigma(G_{\mathrm{ss}})$. Let

$$\mathcal{S}_\chi(G) := \varinjlim L_2^\sigma(G//U)_\chi$$

where U runs through the set of compact open subgroups with $U \cap C(G) \subseteq \ker \chi$. The same estimates as for $\mathcal{S}(G_{\text{ss}})$ show that the convolution on $\mathcal{H}_\chi(G)$ extends to a bounded multiplication on $\mathcal{S}_\chi(G)$.

Consider the map

$$\rho: \mathcal{H}(G) \rightarrow \mathcal{H}_\chi(G), \quad \rho f(g) := \int_{C(G)} \chi(z) f(zg) dz.$$

For appropriately normalised Haar measures, this is a surjective, bounded algebra homomorphism; that is, $\mathcal{H}_\chi(G)$ is a quotient algebra of $\mathcal{H}(G)$. Using ρ , we can pull back $\mathcal{H}_\chi(G)$ -modules to $\mathcal{H}(G)$ -modules. This construction maps essential modules again to essential modules (ρ is a proper morphism in the notation of [17]). Thus we have got a functor $\rho^*: \text{Mod}(\mathcal{H}_\chi(G)) \rightarrow \text{Mod}(\mathcal{H}(G))$. Since ρ is surjective, ρ^* is fully faithful. Thus $\text{Mod}(\mathcal{H}_\chi(G))$ becomes a full subcategory of $\text{Mod}(G)$. It is easy to identify this subcategory with $\text{Mod}_\chi(G)$. If $(V, \pi) \in \text{Mod}_\chi(G)$, then V becomes an essential $\mathcal{H}_\chi(G)$ -module by

$$\pi(f) := \int_{G_{\text{ss}}} f(g) \pi(g) dg.$$

This is well-defined because $f(gz) \pi(gz) = f(g) \pi(g)$ for all $g \in G, z \in C(G)$.

We can extend ρ to a bounded algebra homomorphism $\rho^S: \mathcal{S}(G) \rightarrow \mathcal{S}_\chi(G)$. The map ρ^S has a bounded linear section and its kernel is the closure of $\ker \rho \subseteq \mathcal{H}(G)$. Therefore, bounded algebra homomorphisms $\mathcal{S}_\chi(G) \rightarrow \text{End}(V)$ correspond to bounded algebra homomorphisms $\mathcal{S}(G) \rightarrow \text{End}(V)$ whose restriction to $\mathcal{H}(G)$ vanishes on $\ker \rho$. Equivalently, $\text{Mod}(\mathcal{S}_\chi(G)) \cong \text{Mod}(\mathcal{S}(G)) \cap \text{Mod}_\chi(G)$. Thus $\mathcal{H}_\chi(G)$ and $\mathcal{S}_\chi(G)$ have the required properties.

Theorem 28. *The embedding $\mathcal{H}_\chi(G) \rightarrow \mathcal{S}_\chi(G)$ is isocohomological.*

Proof. Let $\mathcal{H}_\chi(G)^{\text{op}}$ be the opposite algebra of $\mathcal{H}_\chi(G)$, so that $\text{Mod}(\mathcal{H}_\chi(G)^{\text{op}})$ is the category of *right* $\mathcal{H}_\chi(G)$ -modules. Since $\mathcal{H}_\chi(G)^{\text{op}} \cong \mathcal{H}_{\chi^{-1}}(G)$, we have an isomorphism of categories $\text{Mod}(\mathcal{H}_\chi(G)^{\text{op}}) \cong \text{Mod}_{\chi^{-1}}(G)$.

Equip $X \in \text{Mod}(\mathcal{H}_\chi(G)^{\text{op}})$, $V \in \text{Mod}(\mathcal{H}_\chi(G))$ with the associated representations of G . We equip $X \hat{\otimes} V$ with the diagonal representation. Since χ and χ^{-1} cancel, $C(G)$ acts trivially on $X \hat{\otimes} V$. Thus we obtain a bifunctor

$$(20) \quad \text{Mod}(\mathcal{H}_\chi(G)^{\text{op}}) \times \text{Mod}(\mathcal{H}_\chi(G)) \rightarrow \text{Mod}(G_{\text{ss}}), \quad (X, V) \mapsto X \hat{\otimes} V.$$

This functor is evidently exact for extensions with a bounded linear section. Moreover, we claim that $X \hat{\otimes} V$ is projective if X or V are projective. It suffices to treat the case where X is projective. We may even assume that X is a free essential module $X_0 \hat{\otimes} \mathcal{H}_\chi(G)$. The diagonal representation on $X_0 \hat{\otimes} \mathcal{H}_\chi(G) \hat{\otimes} Y$ is isomorphic to the regular representation $\rho_g \otimes 1 \otimes 1$ on $\mathcal{H}(G_{\text{ss}}) \hat{\otimes} X_0 \hat{\otimes} Y$. The intertwining operator is given by $\Phi(x \otimes f \otimes y)(g) := f(g)x \otimes gy$ for all $g \in G$; this function only depends on the class of g in G_{ss} .

Let $X \in \text{Mod}(\mathcal{H}_\chi(G)^{\text{op}})$, $V \in \text{Mod}(\mathcal{H}_\chi(G))$. Then $X \hat{\otimes}_{\mathcal{H}_\chi(G)} V$ is defined as the quotient of $X \hat{\otimes} V$ by the closed linear span of $x * f \otimes v - x \otimes f * v$ for $x \in X$, $f \in \mathcal{H}_\chi(G)$, $v \in V$. Since $\rho: \mathcal{H}(G) \rightarrow \mathcal{H}_\chi(G)$ is surjective, this is the same as $X \hat{\otimes}_{\mathcal{H}(G)} V$, which we have identified with $X \hat{\otimes}_G V$ in Section 3. Thus

$$X \hat{\otimes}_{\mathcal{H}_\chi(G)} V \cong \mathbb{C} \hat{\otimes}_{G_{\text{ss}}} (X \hat{\otimes} V).$$

The same assertion holds for the total derived functors because the bifunctor in (20) is exact and preserves projectives. Especially, we get

$$\mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(G)}^{\mathbb{L}} \mathcal{S}_\chi(G) \cong \mathbb{C} \hat{\otimes}_{G_{\text{ss}}}^{\mathbb{L}} (\mathcal{S}_\chi(G) \hat{\otimes} \mathcal{S}_\chi(G)).$$

Here $\mathcal{S}_\chi(G) \hat{\otimes} \mathcal{S}_\chi(G)$ is equipped with the inner conjugation action of G_{ss} . We identify $\mathcal{S}_\chi(G) \hat{\otimes} \mathcal{S}_\chi(G) \cong \mathcal{S}_{\chi \times \chi}(G \times G)$ as in Lemma 2. By [17, Theorem 35.2], the embedding $\mathcal{H}_\chi(G) \rightarrow \mathcal{S}_\chi(G)$ is isocohomological if and only if $\mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(G)}^{\mathbb{L}} \mathcal{S}_\chi(G)$ is a resolution of $\mathcal{S}_\chi(G)$. Thus the assertion that we have to prove is equivalent to

$$\mathbb{C} \hat{\otimes}_{G_{\text{ss}}}^{\mathbb{L}} \mathcal{S}_{\chi \times \chi}(G \times G) \cong \mathcal{S}_\chi(G).$$

We already know $\mathbb{C} \hat{\otimes}_{G_{\text{ss}}}^{\mathbb{L}} \mathcal{S}(G_{\text{ss}} \times G_{\text{ss}}) \cong \mathcal{S}(G_{\text{ss}})$ because $\mathcal{S}(G_{\text{ss}})$ is isocohomological (Theorem 22) and this condition is equivalent to $\mathcal{S}(G_{\text{ss}})$ being isocohomological.

Now we choose a continuous section $s: G_{\text{ss}} \rightarrow G$; this is possible because G is totally disconnected. It yields bornological isomorphisms

$$\begin{aligned} \Psi': \mathcal{S}_\chi(G) &\rightarrow \mathcal{S}(G_{\text{ss}}), & \Psi' f(g) &:= f \circ s(g), \\ \Psi: \mathcal{S}_{\chi \times \chi}(G \times G) &\rightarrow \mathcal{S}(G_{\text{ss}} \times G_{\text{ss}}), & \Psi f(g, h) &:= f(s(g), s(g)^{-1}s(gh)). \end{aligned}$$

The isomorphism Ψ intertwines the inner conjugation actions of G_{ss} on $\mathcal{S}_{\chi \times \chi}(G \times G)$ and $\mathcal{S}(G_{\text{ss}} \times G_{\text{ss}})$. Thus we get isomorphisms

$$(21) \quad \mathbb{C} \hat{\otimes}_{G_{\text{ss}}}^{\mathbb{L}} \mathcal{S}_{\chi \times \chi}(G \times G) \cong \mathbb{C} \hat{\otimes}_{G_{\text{ss}}}^{\mathbb{L}} \mathcal{S}(G_{\text{ss}} \times G_{\text{ss}}) \cong \mathcal{S}(G_{\text{ss}}) \cong \mathcal{S}_\chi(G).$$

It is easy to see that the composite isomorphism is induced by the convolution map in $\mathcal{S}_\chi(G)$. $\square$

7. APPLICATIONS TO REPRESENTATION THEORY

Let G be a reductive p -adic group, let $C(G)$ be its centre, and let $G_{\text{ss}} := G/C(G)$. So far we have used very large projective $\mathcal{H}(G)$ -module resolutions, which offer great flexibility for writing down contracting homotopies. Now we consider much smaller projective resolutions, which are useful for explicit calculations. We write $\text{Mod}_{(\chi)}(G)$ if it makes no difference whether we work in $\text{Mod}(G)$ or $\text{Mod}_\chi(G)$ for some character $\chi: C(G) \rightarrow \text{U}(1)$. Similarly, we write $\mathcal{H}_{(\chi)}(G)$ and $\mathcal{S}_{(\chi)}(G)$. The actual applications of our main theorem are contained in Sections 7.1, 7.7 and 7.8. The other subsections contain small variations on known results. Our presentation differs somewhat from the accounts in [20, 25] because we want to exhibit connections with K-theory and assembly maps.

7.1. Cohomological dimension. Let $\text{rk } G = \dim \mathcal{BT}(G)$ be the *rank* of G .

Theorem 29. *The cohomological dimensions of the exact categories $\text{Mod}(G)$ and $\text{Mod}(\mathcal{S}(G))$ are (at most) $\text{rk } G$; that is, any object has a projective resolution of length $\text{rk } G$. Similarly, the cohomological dimensions of $\text{Mod}_\chi(G)$ and $\text{Mod}_\chi(\mathcal{S}(G))$ for a character $\chi: C(G) \rightarrow \text{U}(1)$ are at most $\text{rk } G_{\text{ss}}$.*

Proof. The assertions are well-known for $\text{Mod}(G)$ and $\text{Mod}_\chi(G)$. For the proof, equip $\mathcal{BT} = \mathcal{BT}(G)$ with a CW-complex structure for which G acts by cellular maps. Then the cellular chain complex $C_\bullet(\mathcal{BT})$ is a projective $\mathcal{H}(G)$ -module resolution of the trivial representation of length $\text{rk } G$. The chain complex $C_\bullet(\mathcal{BT}) \hat{\otimes} V$ with the diagonal representation of G is a projective $\mathcal{H}(G)$ -module resolution of V for arbitrary $V \in \text{Mod}(G)$. If $V \in \text{Mod}_\chi(G)$, we use the building $\mathcal{BT}(G_{\text{ss}})$ instead; $C_\bullet(\mathcal{BT}(G_{\text{ss}})) \hat{\otimes} V$ is a projective resolution of V in $\text{Mod}_\chi(G)$.

What is new is that we get the same assertions for $\text{Mod}(\mathcal{S}(G))$ and $\text{Mod}(\mathcal{S}_\chi(G))$. Since the argument is the same in both cases, we only write it down for $\mathcal{S}(G)$. Let $V \in \text{Mod}(\mathcal{S}(G))$ and let $P_\bullet \rightarrow V$ be a projective resolution in $\text{Mod}(G)$ of length $\text{rk } G$. Then $\mathcal{S}(G) \hat{\otimes}_G P_\bullet$ has the homotopy type of

$$(22) \quad \mathcal{S}(G) \hat{\otimes}_G^{\mathbb{L}} V \cong \mathcal{S}(G) \hat{\otimes}_{\mathcal{S}(G)}^{\mathbb{L}} V \cong V$$

because $\mathcal{S}(G)$ is isocohomological (Theorem 22); here we use one of the equivalent characterisations of isocohomological embeddings listed in Section 4.2. Equation (22) means that $\mathcal{S}(G) \hat{\otimes}_G P_\bullet$ is a resolution of V . This resolution is projective and has length $\mathrm{rk} G$. $\square$

Conversely, there is $V \in \mathbf{Mod}(G)$ with $\mathrm{Ext}_G^{\mathrm{rk} G}(V, V) \neq 0$. Hence the cohomological dimension of $\mathbf{Mod}(G)$ is equal to $\mathrm{rk} G$. We can even take V tempered and irreducible. Hence $\mathrm{Ext}_{\mathcal{S}(G)}^{\mathrm{rk} G}(V, V) \neq 0$ as well because $\mathcal{S}(G)$ is isocohomological. Thus $\mathbf{Mod}(\mathcal{S}(G))$ also has cohomological dimension equal to $\mathrm{rk} G$. Similarly, the cohomological dimension of $\mathbf{Mod}_\chi(G)$ and $\mathbf{Mod}_\chi(\mathcal{S}(G))$ is equal to $\mathrm{rk} G_{\mathrm{ss}}$.

7.2. Finite projective resolutions. We use a result of Joseph Bernstein to attach an Euler characteristic $\mathrm{Eul}(V)$ in $K_0(\mathcal{H}_{(\chi)}(G))$ to a finitely generated representation $V \in \mathbf{Mod}_{(\chi)}(G)$.

Definition 30. A smooth representation V is called *finitely generated* if there exist finitely many elements $v_1, \dots, v_n$ such that the map

$$\mathcal{H}(G)^n \rightarrow V, \quad (f_1, \dots, f_n) \mapsto \sum_{j=1}^n f_j * v_j$$

is a bornological quotient map.

An admissible representation is finitely generated if and only if it has *finite length*, that is, it has a Jordan-Hölder series of finite length.

Since $\mathcal{H}(G)^n$ carries the fine bornology, the same is true for its quotients. Hence a finitely generated representation necessarily belongs to $\mathbf{Mod}_{\mathrm{alg}}(G)$. In the situation of Definition 30, there exists $U \in \mathbf{CO}(G)$ fixing v_j for all $j \in \{1, \dots, n\}$. Thus we get a bornological quotient map $\mathcal{H}(G/U)^n \twoheadrightarrow V$. Conversely, $\mathcal{H}(G/U)^n$ is finitely generated and projective. Thus a smooth representation is finitely generated if and only if it is a quotient of $\mathcal{H}(G/U)^n$ for some $U \in \mathbf{CO}(G)$. If $V \in \mathbf{Mod}_\chi(G)$, then we may replace $\mathcal{H}(G/U)^n$ by $\mathcal{H}_\chi(G/U)^n$ for some $U \in \mathbf{CO}(G)$ with $\chi|_{U \cap C(G)} = 1$.

Definition 31. An object of $\mathbf{Mod}_{(\chi)}(G)$ has type (FP) if it admits a resolution of finite length by finitely generated projective objects of $\mathbf{Mod}_{(\chi)}(G)$. Such a resolution is called a *finite projective resolution*.

Theorem 32 (Joseph Bernstein). *An object of $\mathbf{Mod}_{(\chi)}(G)$ has type (FP) if and only if it is finitely generated.*

Proof. It is trivial that representations of type (FP) are finitely generated. Conversely, if V is finitely generated, then V is a quotient of a finitely generated projective representation, say, $\partial_0 : \mathcal{H}_{(\chi)}(G/U)^n \twoheadrightarrow V$. By [3, Remark 3.12], subrepresentations of finitely generated representations are again finitely generated. Especially, $\ker \partial_0$ is finitely generated. By induction, we get a resolution (P_n, ∂_n) of V by finitely generated projective objects. By Theorem 29, the kernel of $\partial_n : P_n \rightarrow P_{n-1}$ is projective for sufficiently large n . Hence

$$0 \rightarrow \ker \partial_n \rightarrow P_n \rightarrow \dots \rightarrow P_0 \rightarrow V$$

is a finite projective resolution. $\square$

The *algebraic K-theory* $K_0(\mathcal{H}_{(\chi)}(G))$ is the Grothendieck group of the monoid of finitely generated projective $\mathcal{H}_{(\chi)}(G)$ -modules. This is so because $\mathcal{H}_{(\chi)}(G)$ is a union of unital subalgebras.

Definition 33. Let $V \in \mathbf{Mod}_{(\chi)}(G)$ be finitely generated. Then V is of type (FP) by Bernstein's Theorem 32. Choose a finite projective resolution

$$0 \rightarrow P_n \rightarrow \dots \rightarrow P_0 \rightarrow V \rightarrow 0.$$

The *Euler characteristic* of V is defined by

$$\text{Eul}(V) := \sum_{j=0}^n (-1)^j [P_j] \in K_0(\mathcal{H}_{(\chi)}(G)).$$

We check that this does not depend on the resolution (see also [19, Section 1.7]). Define the Euler characteristic $\text{Eul}(P_\bullet)$ for finite projective complexes in the obvious fashion. Let $P_\bullet$ and $P'_\bullet$ be two finite projective resolutions of V . The identity map on V lifts to a chain homotopy equivalence $f: P_\bullet \rightarrow P'_\bullet$. Hence the mapping cone C_f of f is contractible. The Euler characteristic vanishes for contractible complexes. Hence $\text{Eul}(C_f) = 0$. This is equivalent to $\text{Eul}(P_\bullet) = \text{Eul}(P'_\bullet)$.

Definition 34. Let

$$\text{HH}_0(\mathcal{H}_{(\chi)}(G)) := \mathcal{H}_{(\chi)}(G) / [\mathcal{H}_{(\chi)}(G), \mathcal{H}_{(\chi)}(G)].$$

The *universal trace* is a map

$$\text{tr}_{\text{univ}}: K_0(\mathcal{H}_{(\chi)}(G)) \rightarrow \text{HH}_0(\mathcal{H}_{(\chi)}(G)).$$

If $(p_{ij}) \in M_n(\mathcal{H}_{(\chi)}(G))$ is an idempotent with $\mathcal{H}_{(\chi)}(G)^n \cdot (p_{ij}) \cong V$, then we have $\text{tr}_{\text{univ}}[V] = [\sum p_{ii}]$.

The above definitions are inspired by constructions of Hyman Bass in [2], where $\text{tr}_{\text{univ}} \text{Eul}(V)$ is constructed for modules of type (FP) over unital algebras.

7.3. Traces from admissible representations. Let $W \in \text{Mod}_{(\chi)}(G)$ be an admissible representation. Its integrated form is an algebra homomorphism ρ from $\mathcal{H}_{(\chi)}(G)$ to the algebra $\text{End}_{\text{fin}}(W) := \tilde{W} \otimes W$ of smooth finite rank operators on W . Here $\tilde{W}$ denotes the contragradient representation and $\text{End}_{\text{fin}}(W)$ carries the fine bornology. Composing ρ with the standard trace on $\text{End}_{\text{fin}}(W)$, we get a trace $\text{tr}_W: \text{HH}_0(\mathcal{H}_{(\chi)}(G)) \rightarrow \mathbb{C}$ and a functional

$$\tau_W: K_0(\mathcal{H}_{(\chi)}(G)) \rightarrow \mathbb{Z}.$$

The following computation of τ_W is a variant of [2, Proposition 4.2].

Proposition 35. *Let $V, W \in \text{Mod}_{(\chi)}(G)$, let V be finitely generated projective and let W be admissible. Then $\tau_W[V] = \dim \text{Hom}_G(V, W)$ and $\text{Hom}_G(V, W)$ is finite-dimensional.*

Proof. The functoriality of K_0 for the homomorphism $\mathcal{H}_{(\chi)}(G) \rightarrow \text{End}_{\text{fin}}(W)$ maps $[V]$ to the class of the finitely generated projective module

$$V' := \text{End}_{\text{fin}}(W) \hat{\otimes}_{\mathcal{H}_{(\chi)}(G)} V \cong W \otimes (\tilde{W} \otimes_G V) \cong W^{\dim \tilde{W} \otimes_G V}$$

over $\text{End}_{\text{fin}}(W)$. Thus $\tau_W[V] = \dim \tilde{W} \otimes_G V$. By adjoint associativity,

$$\text{Hom}(\tilde{W} \otimes_G V, \mathbb{C}) \cong \text{Hom}_G(V, \text{Hom}(\tilde{W}, \mathbb{C})) \cong \text{Hom}_G(V, \tilde{\tilde{W}}).$$

We have $W \cong \tilde{\tilde{W}}$ because W is admissible. Thus $\tau_W[V] = \dim \text{Hom}_G(V, W)$. $\square$

Let $V, W \in \text{Mod}_{(\chi)}(G)$, let W be admissible and let V be finitely generated. By Bernstein's Theorem 32, there is a finite projective resolution $P_\bullet \rightarrow V$ in $\text{Mod}_{(\chi)}(G)$. By Proposition 35, $\text{Hom}_G(P_\bullet, W)$ is a chain complex of finite-dimensional vector spaces. Hence its homology $\text{Ext}_{\mathcal{H}_{(\chi)}(G)}^n(V, W)$ is finite-dimensional as well and

$$\begin{aligned} (23) \quad \sum_{n=0}^{\infty} (-1)^n \dim \text{Ext}_{\mathcal{H}_{(\chi)}(G)}^n(V, W) &= \sum_{n=0}^{\infty} (-1)^n \dim \text{Hom}_G(P_n, W) \\ &= \sum_{n=0}^{\infty} (-1)^n \tau_W[P_n] = \tau_W(\text{Eul}(V)). \end{aligned}$$

We call this the *Euler-Poincaré characteristic* $\text{EP}_{(\chi)}(V, W)$ of V and W (compare [20, page 135]).

7.4. Formal dimensions. Evaluation at $1 \in G$ is a trace on $\mathcal{H}_{(\chi)}(G)$, that is, a linear functional $\tau_1: \text{HH}_0(\mathcal{H}_{(\chi)}(G)) \rightarrow \mathbb{C}$. The functional

$$\dim := \tau_1 \circ \text{tr}_{\text{univ}}: K_0(\mathcal{H}_{(\chi)}(G)) \rightarrow \mathbb{C}$$

computes the *formal dimension* for finitely generated projective $\mathcal{H}_{(\chi)}(G)$ -modules. Recall that an irreducible representation in $\text{Mod}_{\chi}(G)$ is projective if and only if it is supercuspidal. Unless $C(G)$ is compact, $\text{Mod}(G)$ has no irreducible projective objects.

We can also define the formal dimension for representations that are square-integrable (see [22]). Let $(V, \pi) \in \text{Mod}_{\chi}(G)$ be irreducible and square-integrable (or, more precisely, square-integrable *modulo the centre* $C(G)$). Since irreducible representations are admissible, V carries the fine bornology. Moreover, square-integrable representations are tempered. Thus the integrated form of π extends to a bounded homomorphism $\pi: \mathcal{S}_{\chi}(G) \rightarrow \text{End}_{\text{fin}}(V)$. Since V is irreducible, this homomorphism is surjective. The crucial property of irreducible square-integrable representation is that there is an ideal $I \subseteq \mathcal{S}_{\chi}(G)$ such that $I \oplus \ker \pi \cong \mathcal{S}_{\chi}(G)$. Thus $\pi|_I: I \rightarrow \text{End}_{\text{fin}}(V)$ is an algebra isomorphism. It is necessarily a bornological isomorphism because it is bounded and $\text{End}_{\text{fin}}(V)$ carries the fine bornology.

Proposition 36. *Let $V \in \text{Mod}_{\chi}(G)$ be irreducible and square-integrable. Then V is both projective and injective as an object of $\text{Mod}_{\chi}(\mathcal{S}(G))$.*

Proof. The direct-sum decomposition $\mathcal{S}_{\chi}(G) \cong \ker \pi \oplus \text{End}_{\text{fin}}(V)$ gives rise to an equivalence of exact categories

$$\text{Mod}_{\chi}(\mathcal{S}(G)) \cong \text{Mod}(\ker \pi) \times \text{Mod}(\text{End}_{\text{fin}}(V)).$$

The representation V belongs to the second factor. The algebra $\text{End}_{\text{fin}}(V)$ is canonically Morita equivalent to $\mathbb{C}$, so that $\text{Mod}(\text{End}_{\text{fin}}(V))$ and $\text{Mod}(\mathbb{C})$ are equivalent exact categories. The easiest way to get this Morita equivalence uses a basis in V to identify $\text{End}_{\text{fin}}(V) \cong \bigcup_{n=1}^{\infty} M_n(\mathbb{C})$. Since any extension in $\text{Mod}(\mathbb{C})$ splits, any object of $\text{Mod}(\mathbb{C})$ is both injective and projective. $\square$

We can also define $K_0(\mathcal{S}_{(\chi)}(G))$, $\text{HH}_0(\mathcal{S}_{(\chi)}(G))$, and

$$\text{tr}_{\text{univ}}: K_0(\mathcal{S}_{(\chi)}(G)) \rightarrow \text{HH}_0(\mathcal{S}_{(\chi)}(G)).$$

It is irrelevant for the following whether we divide by the linear or closed linear span of the commutators in the definition of $\text{HH}_0(\mathcal{S}_{(\chi)}(G))$. The trace τ_1 extends to a bounded trace $\tau_1^{\mathcal{S}}: \text{HH}_0(\mathcal{S}_{(\chi)}(G)) \rightarrow \mathbb{C}$. This induces a functional $\dim^{\mathcal{S}}: K_0(\mathcal{S}_{(\chi)}(G)) \rightarrow \mathbb{Z}$. An irreducible square-integrable representation V defines a class $[V] \in K_0(\mathcal{S}_{(\chi)}(G))$ by Proposition 36; we define its *formal dimension* by $\dim^{\mathcal{S}} V := \dim^{\mathcal{S}}[V]$.

The embedding $\mathcal{H}_{(\chi)}(G) \rightarrow \mathcal{S}_{(\chi)}(G)$ induces natural maps

$$\begin{aligned} \iota: K_0(\mathcal{H}_{(\chi)}(G)) &\rightarrow K_0(\mathcal{S}_{(\chi)}(G)), & [V] &\mapsto [\mathcal{S}_{(\chi)}(V) \hat{\otimes}_{\mathcal{H}_{(\chi)}(G)} V], \\ \iota: \text{HH}_0(\mathcal{H}_{(\chi)}(G)) &\rightarrow \text{HH}_0(\mathcal{S}_{(\chi)}(G)). & [f] &\mapsto [f]. \end{aligned}$$

These maps are compatible with the universal traces and satisfy $\tau_1^{\mathcal{S}} \circ \iota = \tau_1$ and $\dim^{\mathcal{S}} \circ \iota = \dim$.

It is shown in [25] that $\mathcal{S}_{(\chi)}(G)$ is closed under holomorphic functional calculus in the C^* -algebra $C_{\text{red},(\chi)}^*(G)$. Hence

$$K_0(\mathcal{S}_{(\chi)}(G)) \cong K_0(C_{\text{red},(\chi)}^*(G)).$$

It follows also that any finitely generated projective module V over $\mathcal{S}_{(\chi)}(G)$ is the range of a *self-adjoint* idempotent element in $M_n(\mathcal{S}_{(\chi)}(G))$ for some $n \in \mathbb{N}$. Since the trace $\tau_1^{\mathcal{S}}$ is positive, we get $\dim^{\mathcal{S}} V > 0$ unless $V = 0$.

Yet another notion of formal dimension comes from the theory of von Neumann algebras. The $(\chi$ -*twisted*) *group von Neumann algebra* of G is the closure $N_{(\chi)}(G)$ of $\mathcal{H}_{(\chi)}(G)$ or $\mathcal{S}_{(\chi)}(G)$ in the weak operator topology on $L_2(G)_{(\chi)}$. We may extend $\tau_1^{\mathcal{S}}$ to a positive unbounded trace τ_1^N on $N_{(\chi)}(G)$. Any normal $*$ -representation ρ of $N_{(\chi)}(G)$ on a separable Hilbert space is isomorphic to the left regular representation on the Hilbert space $(L_2(G)_{(\chi)} \bar{\otimes} \ell_2(\mathbb{N})) \cdot p_\rho$ for some projection $p_\rho \in N_{(\chi)}(G) \bar{\otimes} B(\ell_2\mathbb{N})$ where $\bar{\otimes}$ denotes spatial tensor products of Hilbert spaces and von Neumann algebras, respectively; the projection p_ρ is unique up to unitary equivalence. We define the *formal dimension* $\dim^N(\rho)$ to be $\tau_1^N(p_\rho) \in [0, \infty]$; this does not depend on the choice of p_ρ .

By definition, we have $\dim^N(L_2(G)_{(\chi)}^n \cdot e) = \dim^{\mathcal{S}}[e]$ if $e \in M_n(\mathcal{S}_{(\chi)}(G))$ is a *self-adjoint* idempotent. Since $\mathcal{S}_{(\chi)}(G)$ is closed under holomorphic functional calculus in the reduced group C^* -algebra $C_{\text{red},(\chi)}^*(G)$, any idempotent element of $\mathcal{S}_{(\chi)}(G)$ is similar to a self-adjoint idempotent. Hence $\dim^N(L_2(G)_{(\chi)}^n \cdot e) = \dim^{\mathcal{S}}[e]$ holds for any idempotent $e \in M_n(\mathcal{S}_{(\chi)}(G))$. We have

$$L_2(G)_{(\chi)} \hat{\otimes}_{\mathcal{S}_{(\chi)}(G)} \mathcal{S}_{(\chi)}(G)^n \cdot e \cong L_2(G)_{(\chi)}^n \cdot e.$$

Therefore, if V is a finitely generated projective left $\mathcal{S}_{(\chi)}(G)$ -module V , then we may view $L_2(G)_{(\chi)} \hat{\otimes}_{\mathcal{S}_{(\chi)}(G)} V$ as a Hilbert space equipped with a faithful normal $*$ -representation of $N_{(\chi)}(G)$; the resulting representation is uniquely determined up to unitary equivalence because any two self-adjoint idempotents realising V are unitarily equivalent in $\mathcal{S}_{(\chi)}(G)$. The formal dimensions from the Schwartz algebra and the von Neumann algebra are compatible in the following sense:

$$\dim^N(L_2(G)_{(\chi)} \hat{\otimes}_{\mathcal{S}_{(\chi)}(G)} V) \cong \dim^{\mathcal{S}}(V).$$

Thus $\dim^{\mathcal{S}} V$ only depends on the unitary equivalence class of the associated unitary representation $L_2(G)_{(\chi)} \hat{\otimes}_{\mathcal{S}_{(\chi)}(G)} V$.

7.5. Compactly induced representations. Equip $U \in \text{CO}(G)$ with the restriction of the Haar measure from G . The map $i_U^G: \mathcal{H}(U) \rightarrow \mathcal{H}(G)$ that extends functions by 0 outside U is an algebra homomorphism. Hence it induces a map

$$(i_U^G)_!: \text{Rep}(U) \cong K_0(\mathcal{H}(U)) \rightarrow K_0(\mathcal{H}(G)), \quad [V] \mapsto [\mathcal{H}(G) \otimes_{\mathcal{H}(U)} V].$$

This is the standard functoriality of K-theory. We denote it by $(i_U^G)_!$ because this notation is used in [17]. We call representations of the form $(i_U^G)_!(V)$ *compactly induced* because $\mathcal{H}(G) \otimes_{\mathcal{H}(U)} V \cong \text{c-Ind}_U^G(V)$ (see [15]).

Let $U, V \in \text{CO}(G)$ and suppose that $gUg^{-1} \subseteq V$ for some $g \in G$. Then we have $i_U^G = \gamma_g^{-1} \circ i_V^G \circ i_{gUg^{-1}}^V \circ \gamma_g$, where γ_g denotes conjugation by g . One checks that γ_g acts trivially on $K_0(\mathcal{H}(G))$. Hence $(i_U^G)_!$ is the composite of $(i_V^G)_!$ and the map $\text{Rep}(U) \xrightarrow{\cong} \text{Rep}(gUg^{-1}) \rightarrow \text{Rep}(V)$ that is associated to the group homomorphism $U \rightarrow V$, $x \mapsto gxg^{-1}$. Let $\text{Sub}(G)$ be the category whose objects are the compact open subgroups of G and whose morphisms are these special group homomorphisms. We have exhibited that $U \mapsto \text{Rep}(U)$ is a module over this category. The various maps $(i_U^G)_!$ combine to a natural map

$$\varinjlim_{\text{Sub}(G)} \text{Rep}(U) \rightarrow K_0(\mathcal{H}(G)).$$

We call this map the *assembly map* for $K_0(\mathcal{H}(G))$ because it is a variant of the Farrell-Jones assembly map for discrete groups (see [12, Conjecture 3.3]), which is in turn closely related to the Baum-Connes assembly map.

The above definitions carry over to $\mathcal{H}_\chi(G)$ in a straightforward fashion. Let $\text{CO}(G; C(G))$ be the set of $C(G)$ -compact open subgroups of G containing $C(G)$. The projection to G_{ss} identifies $\text{CO}(G; C(G))$ with $\text{CO}(G_{\text{ss}})$. As above, we get algebra homomorphisms $i_U^G: \mathcal{H}_\chi(U) \rightarrow \mathcal{H}_\chi(G)$ for $U \in \text{CO}(G; C(G))$. There is an analogue of the Peter-Weyl theorem for $\mathcal{H}_\chi(U)$; that is, $\mathcal{H}_\chi(U)$ is a direct sum of matrix algebras. Therefore, finitely generated projective modules over $\mathcal{H}_\chi(U)$ are the same as finite-dimensional representations in $\text{Mod}_\chi(U)$. This justifies defining $\text{Rep}_\chi(U) := K_0(\mathcal{H}_\chi(U))$. As above, we can factor $(i_U^G)_!$ through $(i_V^G)_!$ if U is subconjugate to V . The relevant category organising these subconjugations is the category $\text{Sub}(G; C(G))$ whose set of objects is $\text{CO}(G; C(G))$ and whose morphisms are the group homomorphisms $U \rightarrow V$ of the form $x \mapsto gxg^{-1}$ for some $g \in G$. Thus we get an assembly map

$$(24) \quad \varinjlim_{\text{Sub}(G; C(G))} \text{Rep}_\chi(U) \rightarrow K_0(\mathcal{H}_\chi(G)).$$

Let $\Gamma(G_{\text{ss}}, dg) \subseteq \mathbb{R}$ be the subgroup generated by $\text{vol}(U)^{-1}$ for $U \in \text{CO}(G_{\text{ss}})$. Since $\text{vol}(U)/\text{vol}(V) \in \mathbb{N}$ for $V \subseteq U$, this group is already generated by $\text{vol}(U)^{-1}$ for maximal compact subgroups $U \subseteq G_{\text{ss}}$. We have $\Gamma(G_{\text{ss}}, dg) = \alpha\mathbb{Z}$ for some $\alpha > 0$ because there are only finitely many maximal compact subgroups and $\text{vol}(U)/\text{vol}(V) \in \mathbb{Q}$ for all $U, V \in \text{CO}(G_{\text{ss}})$. The number α depends on the choice of the Haar measure, of course. We let $\text{size}(U) := \alpha \text{vol}(U/C(G))$, so that $\text{size}(U)^{-1} \in \mathbb{N}$ for all $U \in \text{CO}(G; C(G))$.

Lemma 37. *Let $U \in \text{CO}(G; C(G))$, and let $W \in \text{Mod}_\chi(U)$ be finite-dimensional. Let $c_W: U \rightarrow \mathbb{C}$ be the character of W . Then $\text{tr}_{\text{univ}}(i_U^G)_![W] \in \text{HH}_0(\mathcal{H}_\chi(G))$ is represented by the function*

$$(25) \quad c_{U,W}^G(g) := \begin{cases} \alpha \text{size}(U)^{-1} \overline{c_W(g)} & \text{for } g \in U, \\ 0 & \text{for } g \notin U. \end{cases}$$

Moreover, $\dim(i_U^G)_![W] = \alpha \text{size}(U)^{-1} \dim(W)$. Thus $\dim x \in \alpha\mathbb{Z}$ for all x in the range of the assembly map (24).

A similar result holds for compactly induced projective objects of $\text{Mod}(G)$.

Proof. Since the universal trace is compatible with the functoriality of K_0 and HH_0 , the first assertion follows if $\text{tr}_{\text{univ}}[W] = \text{vol}(U/C(G))^{-1} \overline{c_W(g)}$ in $\text{HH}_0(\mathcal{H}_\chi(U))$. We briefly recall how this well-known identity is proved. We may assume that W is irreducible. Hence there is an idempotent $p_W \in \mathcal{H}_\chi(U)$ with $W \cong \mathcal{H}_\chi(U)p_W$. Thus $\text{tr}_{\text{univ}}[W] = [p_W]$. We can compute $c_W(g)$ for $g \in G$ as the trace of the finite rank operator $f \mapsto \lambda(g)f * p_W$ on $\mathcal{H}_\chi(U)$. This operator has the integral kernel $(x, y) \mapsto p_W(y^{-1}g^{-1}x)$, so that

$$\overline{c_W(g)} = c_W(g^{-1}) = \int_{U/C(G)} p_W(x^{-1}gx) dx.$$

This implies $[\overline{c_W}] = \int_{U/C(G)} [W] dx = \text{vol}(U/C(G)) [W]$ because conjugation does not change the class in $\text{HH}_0(\mathcal{H}_\chi(U))$. We get the formula for formal dimensions because $\dim x = \text{tr}_{\text{univ}}(x)(1)$ and $c_W(1) = \dim W$. This lies in $\alpha\mathbb{Z}$ by construction of α . By additivity, we get $\dim x \in \alpha\mathbb{Z}$ for all x in the range of the assembly map (24). $\square$

7.6. Explicit finite projective resolutions. Let $V \in \text{Mod}_\chi(G)$ be of finite length. Peter Schneider and Ulrich Stuhler construct an explicit finite projective resolution for such V in [20]. We only sketch the construction very briefly. Let $\mathcal{BT}(G_{\text{ss}})$ be the affine Bruhat-Tits building of G_{ss} . One defines a coefficient system $\gamma_e(V)$ on $\mathcal{BT}(G_{\text{ss}})$, which depends on an auxiliary parameter $e \in \mathbb{N}$ (see [20, Section II.2]); its value on a facet F of $\mathcal{BT}(G_{\text{ss}})$ is the—finite-dimensional—space $\text{Fix}(U_F^e, V)$ for certain $U_F^e \in \text{CO}(G)$. The cellular chain complex $C_\bullet(\mathcal{BT}(G_{\text{ss}}), \gamma_e(V))$ with values in $\gamma_e(V)$ is a resolution of V for sufficiently large e ([20, Theorem II.3.1]). It is a finite projective resolution of V in $\text{Mod}_\chi(G)$ because the stabilisers of facets belong to $\text{CO}(G; C(G))$ and the set of facets is G_{ss} -finite.

Proposition 38. *If $V \in \text{Mod}_\chi(G)$ has finite length, then $\text{Eul}(V)$ belongs to the range of the assembly map (24). Hence $\dim(\text{Eul}(V)) \in \alpha\mathbb{Z}$.*

Define the Euler-Poincaré function $f_{\text{EP}}^V \in \mathcal{H}_\chi(G)$ of V as in [20, page 135]. Then $[f_{\text{EP}}^V] = \text{tr}_{\text{univ}} \text{Eul}(V) \in \text{HH}_0(\mathcal{H}_\chi(G))$. Thus $\dim(\text{Eul}(V)) = f_{\text{EP}}(1)$ and $\text{EP}_\chi(V, W) = \text{tr}_W(f_{\text{EP}}^V)$ for all admissible $W \in \text{Mod}_\chi(G)$.

See also [20, Proposition III.4.22] and [20, Proposition III.4.1].

Proof. The finite projective resolution $C_\bullet(\mathcal{BT}(G_{\text{ss}}), \gamma_e(V))$ is explicitly built out of compactly induced representations. Hence $\text{Eul}(V)$ belongs to the range of the assembly map. Lemma 37 yields $\dim(\text{Eul}(V)) \in \alpha\mathbb{Z}$ and allows us to compute $\text{tr}_{\text{univ}} \text{Eul}(V)$. Inspection shows that this is exactly $[f_{\text{EP}}^V]$. $\square$

Proposition 38 yields $\dim V = \dim \text{Eul}(V) \in \alpha\mathbb{Z}$ if V is irreducible supercuspidal. This rationality result is due to Marie-France Vignéras ([25]).

7.7. Euler characteristics and formal dimensions for square-integrable representations.

Theorem 39. *Let $V \in \text{Mod}_\chi(G)$ be irreducible and square-integrable. Let*

$$\iota: K_0(\mathcal{H}_\chi(G)) \rightarrow K_0(\mathcal{S}_\chi(G))$$

be induced by the embedding $\mathcal{H}_\chi(G) \rightarrow \mathcal{S}_\chi(G)$. Then $\iota(\text{Eul}(V)) = [V]$. Hence $[V]$ lies in the range of the assembly map

$$\varinjlim_{\text{Sub}(G; C(G))} \text{Rep}_\chi(U) \rightarrow K_0(\mathcal{H}_\chi(G)) \rightarrow K_0(\mathcal{S}_\chi(G))$$

and $f_{\text{EP}}^V(1) = \dim(\text{Eul}(V)) = \dim^S(V)$. This number belongs to $\alpha \cdot \mathbb{N}_{\geq 1}$ with α as in Lemma 37.

Proof. Choose $e \in \mathbb{N}$ large enough such that $C_\bullet(\mathcal{BT}(G_{\text{ss}}), \gamma_e(V))$ is a projective $\mathcal{H}_\chi(G)$ -module resolution of V . Then

$$\iota \text{Eul}(V) = \sum_{n=0}^{\infty} (-1)^n [\mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(G)} C_n(\mathcal{BT}(G_{\text{ss}}), \gamma_e(V))].$$

Since $\mathcal{S}_\chi(G)$ is isocohomological (Theorem 28), $\mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(G)}^{\mathbb{L}} V \cong V$. Therefore, $\mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(G)} C_\bullet(\mathcal{BT}(G_{\text{ss}}), \gamma_e(V))$ is still a projective $\mathcal{S}_\chi(G)$ -module resolution of V . Since V is projective as well (Proposition 36), this resolution splits by bounded $\mathcal{S}_\chi(G)$ -module homomorphisms. This implies $[V] = \iota \text{Eul}(V)$. The remaining assertions now follow from Proposition 38 and $\dim^S(V) > 0$. $\square$

Theorem 40. *Let $U \in \text{CO}(G; C(G))$ and let $W \in \text{Mod}_\chi(U)$ be finite-dimensional. Then there are at most $\dim(W) \cdot \text{size}(U)^{-1}$ different irreducible square-integrable representations whose restriction to U contains the representation W .*

Proof. Let $V_1, \dots, V_N$ be pairwise non-isomorphic irreducible square-integrable representations whose restriction to U contains W . Let $X := \mathcal{S}_\chi(G) \hat{\otimes}_{\mathcal{H}_\chi(U)} W$. There are natural adjoint associativity isomorphisms

$$\mathrm{Hom}_{\mathcal{H}_\chi(U)}(W, V_j) \cong \mathrm{Hom}_{\mathcal{S}_\chi(G)}(X, V_j)$$

for all j (see [17]). Thus we get non-zero maps $X \rightarrow V_j$. They are surjective and admit bounded linear sections because the representations V_j are irreducible and carry the fine bornology; since the representations V_j are projective (Proposition 36), they even admit G -equivariant bounded linear sections. Thus $V_1, \dots, V_N$ are direct summands of X . Since they are not isomorphic, $\bigoplus V_j$ is a direct summand of X as well. Therefore, $\sum_{j=1}^N \dim^S(V_j) \leq \dim^S X$. Lemma 37 and Theorem 39 yield $\dim^S X = \alpha \dim(W) \mathrm{size}(U)^{-1}$ and $\dim^S(V_j) \geq \alpha$ for all j . Hence $N \leq \dim(W) \mathrm{size}(U)^{-1}$. $\square$

An irreducible square-integrable representation that is not supercuspidal is a subquotient of a representation that we get by Jacquet induction from a proper Levi subgroup. It is desirable in this situation to compute the formal dimension (and other invariants) of V from its cuspidal data. This gives rise to some rather intricate computations; these are carried out in [1] for representations of $\mathrm{GL}_m(D)$ for a division algebra D .

7.8. Some vanishing results.

Theorem 41. *Let $V, W \in \mathrm{Mod}_\chi(G)$ be irreducible and tempered. If V or W is square-integrable, then $\mathrm{Ext}_G^n(V, W) = 0$ for all $n \geq 1$ and*

$$\mathrm{EP}_\chi(V, W) = \begin{cases} 1 & \text{if } V \cong W; \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Since V and W are tempered and $\mathcal{S}_\chi(G)$ is isocohomological, we have

$$\mathrm{Ext}_{\mathcal{H}_\chi(G)}^n(V, W) \cong \mathrm{Ext}_{\mathcal{S}_\chi(G)}^n(V, W).$$

The latter vanishes for $n \geq 1$ by Proposition 36. For $n = 0$ we are dealing with $\mathrm{Hom}_G(V, W)$, which is computed by Schur's Lemma. $\square$

Theorem 42. *If V is irreducible and tempered but not square-integrable, then $f_{\mathrm{EP}}^V(1) = \dim \mathrm{Eul}(V) = 0$.*

Proof. This follows from the abstract Plancherel Theorem and Theorem 41 as in the proof of [20, Corollary III.4.7]. We merely outline the proof. The abstract Plancherel theorem applied to the type I C^* -algebra $C_{\mathrm{red}, \chi}^*(G)$ yields that $f_{\mathrm{EP}}^V(1)$ is the integral of its Fourier transform $W \mapsto \mathrm{tr}_W(f_{\mathrm{EP}}^V)$ with respect to some measure μ , which is called the Plancherel measure. Here W runs through the tempered irreducible representations in $\mathrm{Mod}_\chi(G)$. Proposition 38 asserts that $\mathrm{tr}_W(f_{\mathrm{EP}}^V) = \mathrm{EP}_\chi(V, W)$.

We have $\mathrm{Ext}_{\mathcal{H}_\chi(G)}^n(V, W) = 0$ for all $n \in \mathbb{N}$ and hence $\mathrm{EP}_\chi(V, W) = 0$ unless V and W have the same infinitesimal character. Since the infinitesimal character is finite-to-one, the support of the function $W \mapsto \mathrm{EP}_\chi(V, W)$ is finite. Hence only atoms of the Plancherel measure μ contribute to the integral

$$f_{\mathrm{EP}}^V(1) = \int \mathrm{EP}_\chi(V, W) d\mu(W).$$

These atoms are exactly the square-integrable representations. Now Theorem 41 yields $f_{\mathrm{EP}}^V(1) = 0$ unless V is square-integrable. In addition, this computation shows that $f_{\mathrm{EP}}^V(1) = \dim^S(V)$ if V is square-integrable (compare Theorem 39). $\square$

REFERENCES

- [1] Anne-Marie Aubert and Roger Plymen, *Plancherel measure for $GL(n, F)$ and $GL(m, D)$: explicit formulas and Bernstein decomposition*, J. Number Theory **112** (2005), 26–66. MR2131140
- [2] Hyman Bass, *Euler characteristics and characters of discrete groups*, Invent. Math. **35** (1976), 155–196. MR0432781 (55 #5764)
- [3] J. N. Bernstein, *Le “centre” de Bernstein*, 1–32, Edited by P. Deligne. MR771671 (86e:22028)
- [4] Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften, vol. 319, Springer-Verlag, Berlin, 1999, ISBN 3-540-64324-9. MR1744486 (2000k:53038)
- [5] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local*, Inst. Hautes Études Sci. Publ. Math. (1972), 5–251. MR0327923 (48 #6265) (French)
- [6] Heath Emerson and Ralf Meyer, *Dualizing the coarse assembly map*, J. Inst. Math. Jussieu (2004), <http://arxiv.org/math.OA/0401227> (to appear).
- [7] Alexander Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc., vol. 16, 1955. MR0075539 (17,763c) (French)
- [8] Henri Hogbe-Nlend, *Complétion, tenseurs et nucléarité en bornologie*, J. Math. Pures Appl. (9) **49** (1970), 193–288. MR0279557 (43 #5279) (French)
- [9] ———, *Bornologies and functional analysis*, North-Holland Mathematics Studies, vol. 26, North-Holland Publishing Co., Amsterdam, 1977, ISBN 0-7204-0712-5. MR0500064 (58 #17774)
- [10] James E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990, ISBN 0-521-37510-X. MR1066460 (92h:20002)
- [11] Bernhard Keller, *Derived categories and their uses*, 1996, pp. 671–701. MR1421815 (98h:18013)
- [12] Wolfgang Lück and Holger Reich, *The Baum-Connes and the Farrell-Jones Conjectures in K- and L-theory*, Preprintreihe SFB 478 **324** (2004), Universität Münster.
- [13] Ralf Meyer, *Analytic cyclic cohomology*, Ph.D. Thesis, Westfälische Wilhelms-Universität Münster, 1999.
- [14] ———, *Bornological versus topological analysis in metrizable spaces*, 249–278. MR2097966
- [15] ———, *Smooth group representations on bornological vector spaces*, Bull. Sci. Math. **128** (2004), 127–166. MR2039113 (2005c:22013) (English, with English and French summaries)
- [16] ———, *Comble groups have group cohomology of polynomial growth* (2004), <http://arxiv.org/math.KT/0410597> (to appear in Q. J. Math.)
- [17] ———, *Embeddings of derived categories of bornological modules* (2004), <http://arxiv.org/math.FA/0410596> (eprint).
- [18] A. Yu. Pirkovskii, *Stably flat completions of universal enveloping algebras* (2003), <http://arxiv.org/math.FA/0311492> (eprint).
- [19] Jonathan Rosenberg, *Algebraic K-theory and its applications*, Graduate Texts in Mathematics, vol. 147, Springer-Verlag, New York, 1994, ISBN 0-387-94248-3. MR1282290 (95e:19001)
- [20] Peter Schneider and Ulrich Stuhler, *Representation theory and sheaves on the Bruhat-Tits building*, Inst. Hautes Études Sci. Publ. Math. (1997), 97–191. MR1471867 (98m:22023)
- [21] P. Schneider and E.-W. Zink, *K-types for the tempered components of a p-adic general linear group*, J. Reine Angew. Math. **517** (1999), 161–208, With an appendix by P. Schneider and U. Stuhler. MR1728541 (2001f:22029)
- [22] Allan J. Silberger, *Introduction to harmonic analysis on reductive p-adic groups*, Mathematical Notes, vol. 23, Princeton University Press, Princeton, N.J., 1979, ISBN 0-691-08246-4, Based on lectures by Harish-Chandra at the Institute for Advanced Study, 1971–1973. MR544991 (81m:22025)
- [23] J. Tits, *Reductive groups over local fields*, 29–69. MR546588 (80h:20064)
- [24] François Trèves, *Topological vector spaces, distributions and kernels*, Academic Press, New York, 1967. MR0225131 (37 #726)
- [25] Marie-France Vignéras, *On formal dimensions for reductive p-adic groups*, 225–266. MR1159104 (93c:22034)
- [26] J.-L. Waldspurger, *La formule de Plancherel pour les groupes p-adiques (d’après Harish-Chandra)*, J. Inst. Math. Jussieu **2** (2003), 235–333. MR1989693 (2004d:22009) (French)

E-mail address: `rameyer@math.uni-muenster.de`

MATHEMATISCHES INSTITUT, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, EINSTEIN-STR. 62, 48149 MÜNSTER, GERMANY